


S
333.7
W5 ec
1976

PLEASE RETURN

STATE DOCUMENTS COLLECTION

MAY 24 1978

MONTANA STATE LIBRARY
930 E. Lynd Ave.
Helena, Montana 59601



an
Environmental Plan
for the
Cascade Planning District
Cascade County, Montana

Cole Waldo & Charity Fechter

Resources Development Internship Program
Western Interstate Commission for Higher Education



This report has been catalogued by the WICHE Library as follows:

Waldo, Cole

An environmental plan for the Cascade
County District of Cascade County, Montana /
Cole Waldo, Charity Fechter. -- Boulder, CO :
Western Interstate Commission for Higher
Education, 1976.
52p. + v.p.

1. Land - MT. 2. Environmental policy-
MT. I. Fechter, Charity. II. Cascade
County (MT) Planning Board. III. Western
Interstate Commission for Higher Education.
Resources Development Internship Program.
IV. Title.

The ideas and opinions expressed in this report are those of the author. They do not necessarily reflect the views of the WICHE Commissioners or WICHE staff.

The Resources Development Internship Program has been financed during 1975 by grants from the Economic Development Administration, the Jessie Smith Noyes Foundation, the National Endowment for the Humanities, the Wyoming Office of Manpower Planning; and by more than one hundred and fifty community agencies throughout the West.

WICHE is an Equal Opportunity Employer.

AN ENVIRONMENTAL PLAN
for the
CASCADE PLANNING DISTRICT
of
CASCADE COUNTY, MONTANA

Prepared
by
Cole Waldo
and
Charity Fechter

Supervisor and Coordinator
William G. Monroe, Planning Director

Research Director
James H. Yeagley, Assistant Planner

Maps, Charts, and Graphs
Robert P. Biffle

Typist and Editor
Patty M. Bullock

September, 1976

OCT 10 1994

cascade county planning board



county courthouse great falls, montana 59401

Telephone (406) 761-6700

August 17, 1976

OPEN LETTER TO THE READERS

OF THE

CASCADE PLANNING DISTRICT ENVIRONMENTAL PLAN

This document is the first of two (2) sections which will, when finished, comprise the Comprehensive Land-use Plan for the Cascade Planning District. This plan should be completed by August 31, 1978.

The Cascade County Planning Board was formed by order of the Board of County Commissioners on January 22, 1974. At that time the Planning Board committed itself to an orderly and open process for developing a comprehensive land-use plan for Cascade County. Since that time the Planning Board has hired a professional staff to put together a land-use inventory and ancillary data to formulate a model from which a plan could be written.

The needs, wants, desires and opinions of the citizens living in the different parts of Cascade County will be incorporated in an on-going process by which the Citizen's Advisory Committees, formed throughout the county, will develop sets of goals and policies reflecting the community's viewpoints as to how the planning process should proceed and how the land around them should be used. In this way the people of Cascade County will have an opportunity to control their own future and life style.

The County Planning Department is charged with the task of implementing these goals and policies, by first writing the comprehensive plan and then developing controls and proposed ordinances. All plans, ordinances, and controls will then go through a public hearing process for final review by the citizens of the County.

Open Letter
August 17, 1976
Page 2

This plan is designed to reflect the policies and goals developed by the Cascade Planning District Citizen's Advisory Committee whose long hours of dedication will provide valuable assistance to the planning staff. This plan therefore, belongs to the citizens of the Cascade Planning District.

Sincerely,


Marcia H. Staigmillers
Chairperson
Cascade County Planning Board

WGM/pb

A C K N O W L E D G E M E N T S

Cascade County Planning Board

Marcia Staigmiller - Chairperson

Malvin Merja - Vice-chairperson

Richard Golie

Franklin Vanatta

Marjorie McCafferty

Dan Thurber

Robert Erickson

Cascade County Commissioners

Edward Shubat - Chairman

John St. Jermain

Lawrence Fasbender

Cascade County Planning Board Staff

William G. Monroe, Planning Director

James H. Yeagley, Assistant Planner

Robert P. Biffle, Cartographer

Patty M. Bullock, Typist and Editor

William Murray, Coordinator, Civil Defense

TABLE OF CONTENTS

ABSTRACT.	page 1
INTRODUCTION	2
DESCRIPTION AND HISTORY	3
SECTION I - LANDSCAPE UNITS	6
Rivers, reservoirs, lakes, and streams	9
Alluvial Lowlands	11
Sedimentary terraces or benches	14
Sedimentary foothills	18
Igneous foothills	21
Igneous uplands	23
Buttes	25
SECTION II - CRITICAL HAZARD AREAS.	28
Flood hazard areas	29
Ice jams	31
Mass movement	32
Earthquakes.	35
SECTION III - PUBLIC OPEN SPACE	37
Part I - Explanation	37
Part II - Defining open space	38
Part III - Need for open space	39
Part IV - Implementation	46
SECTION IV - PRIORITY RESOURCES	48
Agricultural lands	49
Fisheries	50
Minerals and mineral aggregates	51
Water resources	52
APPENDIX A - DEFINITION OF LANDSCAPE UNITS	A-1
APPENDIX B - GEOLOGIC TIME SCALE	B-1
APPENDIX C - ENDANGERED SPECIES	C-1
APPENDIX D - ALTERNATIVE ENERGY	D-1
Solar energy	D-6
Sun angle	D-7
Wind	D-9
Local climatological data	D-11
GLOSSARY	G-1
BIBLIOGRAPHY	BB-1

MAPS, CHARTS, AND GRAPHS

	<u>PAGE</u>
CASCADE PLANNING DISTRICT MAP	5
DEGREES OF TOPOGRAPHIC SLOPE	8
LANDSCAPE UNITS	27
TOPOGRAPHY	36
STREAM FLOW CHART (MISSOURI RIVER MAIN STEM)	54
PRIORITY RESOURCES	55
LANDSCAPE UNITS ON CROSSECTION	A-3
ENERGY RESOURCES COST CHART	D-5
SUN ANGLE AT GREAT FALL, MT.	D-8
ANNUAL ALL WEATHER WIND ROSE	D-10
LOCAL CLIMATOLOGICAL DATA CHARTS	D-12
METEOROLOGICAL DATA	D-13
NORMALS, MEANS, AND EXTREMES	D-14

ABSTRACT

This report is the environmental plan for the Cascade Planning District. It is made up of four (4) basic elements: 1) Landscape Units, their soils, geology, topography, hydrology and biology, with suggested development and conservation policies for each landscape unit; 2) Critical Hazard Areas, defined, with suggested performance standards and policies for each hazard type; 3) Public Open Space Program, with some methods of implementation; 4) Priority Resource Areas, with suggested policies and actions to protect these resources. A glossary of terms that may be unfamiliar is also included. There are, as well, four (4) appendices: 1) Definition of Landscape Units; 2) Geologic Time Scale; 3) Endangered Species; 4) Alternative Energy.

ENVIRONMENTAL PLAN FOR THE CASCADE PLANNING DISTRICT

INTRODUCTION

The comprehensive plan for Cascade County is being developed in two (2) parts: an environmental plan and a community plan for each of the eleven (11) planning districts. The environmental plan deals with the natural setting for the social and economic activities to be covered in the community plan. With the completion of each environmental plan residents within the various planning districts will be contacted, then a community plan will be developed by the Citizen's Advisory Committee in each district.

Each district's environmental plan is made up of four (4) basic elements: landscape units, critical hazard areas, open space program and priority resource areas. Each element performs a specific purpose in incorporating environmental data and policies into the comprehensive plan.

Landscape Units - The county has been broken into nine (9) landscape units, with each representing a particular combination of soils, geology, topography, hydrology and biology. The landscape units therefore provide a simplified framework for development and conservation policies and apply to the same landscape unit throughout the county.

Critical Hazard Areas - Areas of known soil instability, flooding, stream and river bank erosion as well as wind erosion are identified by text. In the interest of protecting life and property, performance standards and policies have been developed for each type of hazard.

Public Open Space Program - Open space designations in this plan are restricted to sites which can be publicly controlled and maintained and which can serve a function for recreation, scenic value, wildlife habitat, historical, cultural or educational interest.

Priority Resource Areas - Those resources with economic value are identified and described. Policies and actions are set forth in the plan to protect forest (silvaculture), agriculture, and mineral resources.

When combined, the elements of the environmental plan provide the foundation for social and economic policies in the community plan. There is established a systematic format for dealing with hazards, economic resources and open space needs, as well as developing relationships between soils, air, water, and living organisms. The plan sets a framework for policies and performance standards but is flexible enough to accomodate changes in policies - should such occur. The policies provide specific guidelines for establishing ordinances and making decisions.

DESCRIPTION AND HISTORY

Cascade County is located in north-central Montana east of the continental divide. Topography of the area varies from very mountainous in its southern third to rolling plains in its northern portions. Elevations range from peaks over 8000 feet to river bottoms near 2700 feet. The county is drained by four major rivers and creeks which include the Missouri River, Sun River, Smith River, and Belt Creek. The climate has many "Continental" characteristics with an important exception being the so called "chinook" which is an occasional warm winter wind. Warm summers and cold arctic winters are normal in the area. Rain fall generally occurs in the spring and summer months.

The first penetration by white men of the area within the present boundaries of Cascade County was made by Lewis and Clark Expedition in June, 1805. The area was soon named for the many falls and cascades which characterize the area.

Organization of the county took place on September 19, 1887, when Cascade County was formed from parts of adjacent Chouteau, Lewis and Clark, and Meagher Counties. 1887 was also the year when railroads first arrived and connected the area to other parts of the nation. Spur lines were constructed to many local mining areas making Great Falls the transportation hub for much of north-central Montana.

Cascade County is also the site of Montana's first hydroelectric plant: Black Eagle Plant. It was built in 1890 in the canyon near Great Falls. Since that date a series of five (5) hydroelectric dams have been constructed on the Missouri River within the county.

Mining and agriculture have played a very important role in development of the area. Major mining operations took place in the southeastern part of the county but have since been discontinued. Smelting operations for copper and zinc have played an important part in the development of the Great Falls area.

Great Falls is a regional center for agricultural exchange and supply. Cattle and grain are the primary agricultural products and provide the economic base for the area.


Another important factor to development of the area was the addition of Malmstrom Air Force Base and its numerous projects. Light diversified manufacturing has also been important to growth of the area as well as increased services available to customers.

Population of Cascade County as of 1970 was 81,631. Approximately 87% is urban, 10% rural non-farm, and 3% rural farms. Distribution of the population closely follows the Missouri and Sun Rivers. Incorporated towns within the county include Great Falls, Cascade, Belt, and Neihart. Many unincorporated towns and villages also exist.

The Cascade Planning District is located in the southwestern corner of the county. It is the largest of the eleven (11) planning districts, occupying 434,342 acres (approximately 679 square miles). At its widest point the district is thirty (30) miles and its length is thirty-four (34) miles.

The district has the greatest variation of topography in the county. Within its borders exist alluvial lowlands, terraces and benches, foothills, buttes and uplands.

Cascade is the only incorporated town in the district and is centrally located along the Missouri River. Many communities can be found in the region and are often closely associated with school districts. The population is primarily found along the Missouri River and within the incorporated town of Cascade.



CASCADE PLANNING DISTRICT



SECTION I

LANDSCAPE UNITS

ENVIRONMENTAL PLAN FOR THE CASCADE PLANNING DISTRICT

SECTION I - LANDSCAPE UNITS

The landscape unit section is the first of four (4) basic elements comprising the environmental plan. This section provides the framework for setting forth environmental policies based on existing natural conditions. The landscape unit designations and the recommendations attached to each unit will be useful in establishing zones and performance standards in the Community Plan.

Each landscape unit reflects a set of characteristics which constitutes a natural process. The soils, hydrology, biology, and landforms are inter-related as a functional, natural unit. Environmental policies are established to reflect the overall natural process of landscape unit, as well as specific resource limitations.

Every parcel of land in the county is within a landscape unit. The landscape units found in the Cascade Planning District are:

1. Rivers, reservoirs, lakes and streams
2. Alluvial lowlands
3. Sedimentary terraces or benches
4. Sedimentary foothills
5. Igneous foothills
6. Igneous uplands
7. Buttes

Other landscape units within Cascade County:

1. Sedimentary uplands
2. Metamorphic uplands

The flexibility of the environmental plan allows the addition of other units, as warranted by specific conditions. Development policies attached to each landscape unit apply to all lands within the landscape unit. As a general rule, no parcel will fall in more than one (1) landscape unit. An individual landowner, however, may have land in two (2) or more units.


For each landscape unit the following information is provided:

1. Definition of the landscape unit.
2. Extent and description of that landscape unit within the planning district.
3. The slope, geology, soils, vegetation, wildlife, and land-use characteristics.
4. Example area.
5. Suggested development policies for that landscape unit.
6. Recommended actions to implement the policies.

A map of the planning district showing the landscape units at a scale of four (4) miles to the inch (1:253,440) accompanies this section. The map is as accurate as possible at this scale and there should be no question as to which landscape unit a particular parcel belongs. In cases where there is some question, an on-site inspection may be required to make a determination. Given the detailed description provided in this section, the reader should have little difficulty in placing a site in the appropriate landscape unit.

Appendix A defines all the landscape units found in Cascade County. Six (6) of the nine (9) landscape units are found in the Cascade Planning District.

DEGREES OF TOPOGRAPHIC SLOPE



RIVERS, RESERVOIRS, LAKES, STREAMS - LANDSCAPE UNIT

DEFINITION - Includes water bodies either flowing or standing all or part of the year. Lands immediately adjacent to water bodies which directly influence the physical, biological, and chemical properties of the water are also included in this landscape unit.

EXTENT AND DESCRIPTION - Lakes, reservoirs and streams in the Cascade Planning District are mainly centered around the Missouri River drainage. Lakes in the district include Schrammeck Lake (Section 7, Township 17 North, Range 2 East) and a number of small lakes on the terrace and benches landscape unit.

Reservoirs are spread throughout the district in all landscape units except the alluvial lowlands and buttes. Major reservoirs include Middle Creek Lake, Hound Creek Reservoir, Rocky Reef Reservoir, and Elizabeth Reservoir.

The Missouri River is the major drainage system of the area. Its tributaries include: Muddy Creek, Antelope Creek, Castner Coulee, Bird Creek, and the Dearborn River. The Hound Creek - Smith River drainage is very important on the eastern side of the Cascade Planning District.

VEGETATION - The natural vegetation is similar to that found in the alluvial lowlands: native cottonwood; willow; brush, including choke-cherry and serviceberry; native grasses and shrubs, such as slough grass, wild barley, sage, rushes, and sedges. The rivers have a greater variety of marsh grasses and algae.

WILDLIFE - Fish and waterfowl are the most important wildlife of this landscape unit. Game fish, such as trout, pike and catfish, can be found in all annual streams and within many of the reservoirs and lakes. Waterfowl includes duck, gulls, and occasional swans and geese. Blackbirds, magpies, and other birds are also found in this area.

GEOLOGIC HAZARDS - Flooding is the major hazard of this landscape unit. Water control by the dam system on the Missouri River south of Cascade County has limited the problem to some extent but high water flow and blockages of the river by ice jams can cause flooding on the lowlands.

CHARACTERISTICS -

GEOLOGY - The geology of the rivers and streams landscape unit in the Cascade Planning District is primarily of the Quaternary Period

(3 million years ago). The rivers flow on dune sands and glacial lake deposits which have gradually filled the valley bottom. Alluvium colluvium can also be found beneath the waters in the southwestern part of the Cascade Planning District where erosion has caused landslides into the river bed. The streams cut through the uplands landscape unit forming the deep canyons characteristic of the area.

SOILS - Soils are descibed more completely under the individual landscape units.

SLOPE - Mostly gently sloping riverbank.

EXAMPLE AREA LOCATION - Schrammeck Lake in Section 7, Township 17 North, Range 2 East. Missouri River in the Chestnut Valley.

SUGGESTED POLICIES -

1. Since the rivers, reservoirs, lakes, and streams are of the highest priority in terms of recreation, aesthetics, agriculture, and wildlife habitat, every effort should be made to preserve their present integrity.
2. Wildlife habitat should be protected from land-use changes.
3. Rechanneling of streams should be discouraged.
4. Public access to all rivers and streams should be provided at regular intervals.
5. Areas designated for their unique and fragile or recreational potentials should be protected by whatever means are most feasible.
6. Vegetation buffer zones along streams should be managed primarily for water quality, stream protection, and wildlife habitat.
7. Efforts should be directed toward maintaining or improving water quality.

RECOMMENDED ACTIONS -

1. The county should coordinate all proposed actions affecting the Missouri River with the Montana Department of Fish and Game and the U.S. Soil Conservation Service to prevent adverse effects on wildlife.
2. Agricultural lands would be given priority, but special efforts should be made to prevent erosion.
3. Establish minimum flows to protect the wildlife, water quality, and recreational uses of this landscape unit.

ALLUVIAL LOWLANDS - LANDSCAPE UNIT

DEFINITION - The plains occupying the valley floors which are the result of deposition of material by water. The floodplain is the major conspicuous feature of this landscape unit.

EXTENT AND DESCRIPTION - Alluvial lowlands in the Cascade Planning District are closely associated with the Missouri River and its tributaries.

Flatlands or floodplains extend in an arc around Cascade from the southwest near Hardy Creek to the northeast near Riverdale on the eastern side of the river. The west bank is higher having few places of alluvial lowlands. Upper Muddy Creek and Rocky Gap Creek in the northern part of the district are also regions of alluvial lowlands. Extension of the alluvial lowlands in this report is to the 3400 foot level.

VEGETATION - Cultivation accounts for the largest percentage of vegetation in the alluvial lowlands. Modification of the natural landscape for agricultural purposes has caused much of the natural vegetation to become scarce.

Natural vegetation includes: native cottonwoods; willow; brush, chokecherry, serviceberry, etc.; native grasses and shrubs, slough grass, wild barley, rush, sedge, and sage.

WILDLIFE - Excellent habitats for big game are found in the alluvial lowlands. Mule deer and whitetail deer can be found year round while antelope use the area as their winter range. Game birds include: pheasant, sharptail grouse, Hungarian partridge and some ducks. Many other birds such as robins, blackbirds and sparrows can also be found in this area. Small game consisting of otter, ground squirrels, rodents and many others also inhabit this landscape unit. Bald Eagle and osprey are sometimes found to use this area as a feeding ground.

GEOLOGIC HAZARDS - The major geologic hazards in the alluvial lowlands of the study area are stream flooding, bank erosion and dune sand formation. Flooding is the primary hazard at the lower elevations along the Missouri River and its tributaries. Development of Holter, Hauser, and Canyon Ferry Dams has regulated water flows along the Missouri in recent years. Flooding on small drainages is evident when high rainfall and snowfall occurs, giving them the potential to be hazardous. Bank erosion is often a result of flooding and can

be hazardous to nearby development. Dune sand formation often occurs in the alluvial lowlands and is generally caused by wind erosion. Sands can be detrimental to agricultural systems in the area.

CHARACTERISTICS -

GEOLOGY - Alluvial lowlands are composed of sands, silt and clay eroded from the local landscape or deposited by glacial lakes. The area is relatively flat with many cut off stream meanders present where the former river bed existed. Gravel lenses composed of subrounded moderately well sorted materials are found in the area. The age of the sediments is believed to be of the early Quaternary period (3 million years ago). The alluvial formations follow the river beds and the adjacent low flatlands or floodplain. These formations lie on the silt-shales of the Marias River Formation. Dune sands have formed on much of the flatlands and are presently active. They are the result of wind erosion of the many sandstones found in the area.

SOILS - The majority of the soils in this landscape unit belong to the Straw-Libra-Nesda association. These soils are mainly deep, well drained, dark colored loams and sandy loams. There are some shallow, very gravelly soils in narrow bands on the floodplains adjacent to streams. Straw soils are deep, well drained, dark colored, stratified, loamy soils. Ponding and flooding hazards limit the use of this soil for building sites.

Libra soils are deep, well drained, dark colored, calcareous soils with fine sandy loam. The surface is made up of loam and clay loam layers. The subsoils are sandy loam. Building sites are limited by the flooding and ponding hazards.

Nesda soils are gravelly and cobbly loams with loose sand and gravel subsoils to a depth of 60 inches. Ponding and flooding hazards limit the use of this soil for building sites. The Straw-Libra-Nesda association is used primarily for small grain, hay, pasture, and grazing. Part of the area in this association is irrigated.

Absher-Nobe association soils are also found in this landscape unit. These soils are deep, very slowly permeable, saline-alkaline clays. The clay content is unfavorable for most uses, as is the salinity and alkalinity. This association is used mainly for grazing.

SLOPE _ Gentle; 0 - 2%.

EXAMPLE AREA LOCATION - Chestnut Valley along the Missouri River.

SUGGESTED POLICIES -

1. Low density activities, such as agriculture, shall be preferred because of the flood hazard.
2. Housing developments will be discouraged, and will not be allowed in designated floodplain. Soils which are prone to high groundwater are recommended for agricultural use only.
3. Hospitals, schools, subdivisions, rest homes, and commercial and industrial developments will be strictly prohibited from the floodplains.
4. All developments shall be reviewed by the appropriate agencies and the Citizen's Advisory Committee for impact on the environment.
5. Dikes, Levees, and other methods of flood control will be considered a land-use problem rather than engineering problem.

RECOMMENDED ACTIONS -

1. Curtail man's activities, except agriculture, on the floodplains and floodways.
2. Map the floodway and floodway fringe for at least the 100 year flood. Ideally, the floodplain should be delineated for the 5, 10, 20, and 50 year floods as well.
3. Land-use controls should reflect the environmental limitations on the alluvial lowlands.
4. Any further development should occur within existing community boundaries only.

SEDIMENTARY TERRACES OR BENCHES - LANDSCAPE UNIT

DEFINITION - The relatively flat or gently sloping topographic surfaces which mark former valley floor levels. Stream downcutting has caused the terraces to stand higher than the present valley floor. Terraces are traditionally composed of alluvial deposits of sand, silt and gravel. Some of the necessary gravels, however, have either not been deposited by the stream or have been subsequently eroded. Therefore, the term "bench" may be more appropriate. There are some remnant terrace sand, silt and gravel deposits throughout the county, so the terms "terrace" and "bench" may be used interchangeably throughout the plan.

EXTENT AND DESCRIPTION - Terraces and benches are the relatively flat areas above the alluvial lowlands. Gentle hills and draws are characteristic of the area. Elevations extend from 3400 feet to 4000 feet. Grazing and dryland farming are the largest uses of the land with some sprinkler irrigation being introduced.

Terraces and benches account for most of the agricultural land in the district and have been developed extensively for that purpose.

VEGETATION - Cultivation accounts for much of the vegetation in this landscape unit. Dryland farming, haylands and grazing are the primary uses and alteration of the natural vegetation is the result.

Native grasses of the area include western wheatgrass, bluegrass, and bluebunch wheatgrass. Few trees are found in the area and brush is limited to draws and coulees.

WILDLIFE - Big game found in this landscape unit includes mule deer, white-tail deer and antelope. Sharptail grouse, pheasant and Hungarian partridge also inhabit the area.

A variety of small animals, such as ground squirrels, also occupy the area.

GEOLOGIC HAZARDS - Major hazards in this landscape unit include water erosion and wind erosion (blowouts). Erosion is particularly evident where the terrace landscape unit meets the alluvial lowlands landscape unit and where water flows are at their highest. Wind averages in this area are nearly the highest in the nation and account for damages from blowouts and dune formation. Slumping on minor scales is evident in some places.

CHARACTERISTICS -

GEOLOGY

- Major formations were deposited in the early stages of the Cretaceous Period (100 million years ago). The Kootenai formation, composed of red, purple, and green siltstone with beds of sandstone and limestone, is the first of the formations. It is located along Hound Creek and Spanish Coulee.

The Blackleaf formation is the next major formation and it is often referred to as the Colorado Shales. It has three (3) major members which are the Flood, Taft Hill and Bootlegger deposits. They are all composed of sandstone, siltstone and shales with some pebble conglomerates. The last major formation is the Marias River Formation which consists of black shales and black sandstones.

Other deposits which can be found in this landscape unit are the dune sands and terrace deposits. The sands are believed to be from nearby formations of sandstone or glacial lake deposits which have been eroded by the wind. Terrace gravel found in this district is composed of well-rounded poorly sorted pebbles and cobbles which are believed to have been deposited during the Ice Age of the Quaternary Period (3 million years ago). Presently these terrace gravels are the major gravel sources for the area.

SOILS

- The Pilgrim-Sinnigam-Rough broken land association is fairly important in this landscape unit. These are well drained, dark colored, clay loam and channery or very stony loam soils 15 to 40 inches deep to sandstone or shale. Steep areas soils are stony. Pilgrim soils have dark colored, clay loam and stony clay loam surface layers. The subsoils are silty clay and are calcareous below depths of 20 to 26 inches. Depth to shale is between 30 to 40 inches. The unfavorable clay content of the Pilgrim soils limit their suitability for most uses.

Sinnigam soils are dark colored and very stony with a depth to hard sandstone of 10 to 20 inches. These well drained soils occur on convex positions in uplands as well as along the bedrock bench edges. Depth to bedrock is the limiting factor for most uses.

Rough broken land soils are deep, very stony loams that occur on steep slopes. The Pilgrim-Sinnigam-Rough broken land association

is used for nonirrigated small grain and hay on the gently sloping and rolling topography. The steep and very stony areas are used for grazing.

Another important association is that of the Winifred-Yawdim-Butchert soils. Winifred soils are moderately dark, silty, clay loam surface soils and calcareous, silty clay or silty clay loam subsoils. The depth to clayey shale in these well drained soils is 20 to 40 inches. Frost heave and clay content make this type soil unsuitable for number of reasons.

The calcareous, light colored, shallow, clayey Yawdim soils are less than 20 inches deep to shale. Depth to bedrock and clay content are the major limitations on most uses.

Butchert soils are channery and stony loam that are less than 10 inches deep to sandstone. The coarse fragments and depth to bedrock put severe limitations on this soil for most uses. This soil association is used primarily for grazing, though some areas are used for small grain and hay.

Azaar-Gollaher-Tally association soils are dark colored, well drained sandy soils. Depth to bedrock and rapid permeability are the major limiting factors. This association is used for grazing and nonirrigated small grain and hay.

The Rough broken land association is dominated by steep land broken by numerous intermittent drainage channels. Runoff is high and geologic erosion is active. Soils of this type are generally used for grazing and wildlife.

The Hanson-Dumco-Faines association is of minor importance in the sedimentary terrace landscape unit. More detail on this association may be found in the sedimentary foothills landscape unit. Absher-Nobe association soils are present but have minor importance. This association is discussed in more detail in the alluvial lowlands landscape unit.

SLOPE - Steep, scarp upslope followed by a rolling plain. Slope is highly variable, gentle (0 - 10%) on the plain, steep (over 25%) on the upslope.

EXAMPLE AREA LOCATION - Section 8, Township 17 North, Range 1 West.

SUGGESTED POLICIES - Because of the variability of limitations, extensive on-site inspection and evaluation should be made by qualified professionals before any proposed action is taken.

1. Because present land uses are predominately agricultural, special consideration should be given to this use.
2. Maintenance of natural vegetation should be a prime consideration in order to preserve the landscape, discourage erosion, and maintain wildlife habitates.

RECOMMENDED ACTIONS

1. Develop land-use controls which reflect the environmental limitations and assets of the terrace.
2. Encourage development in a form which reflects environmental limitations and the natural landscape.

SEDIMENTARY FOOTHILLS - LANDSCAPE UNIT

DEFINITION - The foothills lie between the sedimentary terraces and the igneous uplands. The lower parameter is the 4000 foot contour and the upper parameter, the break between foothills and uplands, is the 5000 foot contour line.

EXTENT AND DESCRIPTION - Foothills extend in a relatively narrow band around the igneous uplands which are in the southwestern part of the Cascade Planning District. Steep rounded hills sloping upward to the uplands are characteristic of the area. Grazing is the primary use of this land.

VEGETATION - Vegetation is similar to that in the terraces or benches landscape unit with a higher percentage being of native origin. Land use is primarily agricultural grazing lands using native vegetation. Haylands can be found in the creek bottoms. Trees are more abundant. Grasses include: western wheatgrass, bluegrass, bluebunch wheatgrass, and numerous shrubs and forbs.

WILDLIFE - Wildlife in the foothills is highly similar to that of benches and terraces. Big game includes: mule deer, whitetail deer, and antelope. The area is sometimes used as winter range for elk. Game birds include grouse, Hungarian partridge, and pheasant. Small animals of a great variety are also present in this landscape unit.

GEOLOGIC HAZARDS - Geologic hazards in this area include faults, landslides, slumping and soil creep. Some erosion is also evident and could be potentially hazardous.

Earthquake faults can be found in the southern part of the Cascade Planning District. The Carter Ranch Fault in Township 16 North, Range 2 East is one of the largest in the area. Several other smaller faults are located in the same area. Landslides are common in this area where the sharp drop-off from the uplands landscape unit is gradually eroded. During high rainfall periods slumping and soil creep occur on some of the steeper foothills where soil water saturation is high.

CHARACTERISTICS -

GEOLOGY - Deposits in this landscape unit are similar to the terrace and bench landscape unit but are more inconsistent in size and

exposures. The Kootenai Formation is the oldest formation in the unit being of the Lower Cretaceous Period (100 million years old). It is composed of highly colored siltstone, sandstones, and limestones.

The Blackleaf formation including the Flood, Taft Hill and Bootlegger members is also found in this unit. Exposures are scattered and inconsistent but are of varying shades of sandstone, siltstone and shale which were probably deposited when the area was at the shores of an inland sea. The Marias River Formation, another black sandstone-shale formation is also evident in this landscape unit. Alluvium colluvium composed of sand, silt, clay and basalt (lava) fragments is another important feature of this landscape unit. Exposures often are found at the edge of the foothills unit, near the uplands landscape unit. Thickness of these deposits is often as much as 50 feet. Faults and folding have played an important part in the geology of this area. Three (3) minor faults and one (1) major fault can be found in this area. The Carter Ranch Fault is a reverse fault in the Disturbed Belt caused by north-south compressional stresses of two (2) tectonic plates. The Carter Ranch Fold is another result of these stresses along with several other forces which have gradually caused deformations of the strata beds.

SOILS

- The Hanson-Dumco-Faines association is the most important in this district. These soils are primarily black, stony loam and loam soils more than 20 inches deep to limestone bedrock. Hanson soils have a black loam or very stony loam surface layer and nearly white, strongly calcareous, very stony loam subsoils. The slope and coarse fragments connected with this soil provide some limits to its suitability for certain purposes.

Dumco soils are similar to Hanson soils, but the depth to bedrock is less than 20 inches. Steepness and depth to bedrock are severely limiting for most uses.

Faines soils are made of black, silt loam surface layers and dark loam, silty clay loam subsoils that are calcareous below depths of 20 to 24 inches. Depth to bedrock and slope limit some uses. These soils are used for nonirrigated small grain and hay on the sloping and rolling topography. Grazing and woodland form the uses on other areas.

Pilgrim-Sinnigam-Rough broken land association is of minor importance and is discussed in greater detail in the sedimentary terrace landscape unit. The Mucet-Sangrey-Rock outcrop association is also of minor importance and is discussed in the igneous upland landscape unit.

SLOPE -Variable, up to 25% or more.

EXAMPLE AREA LOCATION - Area near Harris Mountain below 5100 feet in Section 3, Township 16 North, Range 1 West.

SUGGESTED POLICIES-

1. Grazing and other agricultural activities should be utilized to the fullest.

RECOMMENDED ACTIONS -

1. If any development takes place, extensive on-site evaluations should be made. Criteria weighed should be;
 - a. Soils limitations
 - b. Ground and surface water; depth to groundwater
 - c. Geologic hazards
 - d. Slope
 - e. Adverse effect on wildlife habitat
 - f. Visual impact
 - g. Effects on recreational uses and access
2. Land-use controls should be based on the aforementioned criteria.
3. Agricultural used should be given top priority.

IGNEOUS FOOTHILLS - LANDSCAPE UNIT

DEFINITION - The igneous foothills occupy the area between the sedimentary terraces and the igneous uplands and are adjacent to the sedimentary foothills. The lower parameter is the 4000 foot contour and the upper parameter is the 5000 foot contour.

EXTENT AND DESCRIPTION - The igneous foothills apron the base of the igneous uplands, and are found primarily in the southern portion of the Cascade Planning District. Steep rounded hills slope upward to the uplands. Grazing on native vegetation is the major land use.

VEGETATION - The foothills support a higher percentage of native vegetation than the benches, but haylands are found in some of the creek bottoms. Trees are more abundant in this landscape unit with coniferous varieties dominant. Grasses include: Western wheatgrass, bluegrass, and bluebunch wheatgrass.

WILDLIFE - Wildlife in the foothills is highly similar to that of benches and terraces. Big game includes: mule deer, whitetail deer, and antelope. The area is sometimes used as winter range for elk. Game birds include grouse, Hungarian partridge, and pheasant. Small animals of a wide variety are also present in this landscape unit.

GEOLOGIC HAZARDS - The geologic hazards inherent to this area are faults and landslides. Because of the nature of this area and the igneous origin, there are only isolated instances of soil creep and slumping. Earthquake faults are found in the southern portion of the Cascade Planning District with the largest being the Carter Ranch Fault.

CHARACTERISTICS -

GEOLOGY - This area is composed primarily of the Adel Mountain Volcanics. These potash-rich Late Cretaceous Age (60 million years old) or early Quarternary (20 million years old) volcanics are believed to have intruded before the main diastrophism of the Disturbed Belt.

SOILS - The major soils association is the Mucet-Sangrey-Rock outcrop association which is discussed in the igneous uplands landscape unit. Steepness and depth to bedrock put severe limits on this soil for most uses.

SLOPE - Variable, up to 25%.

EXAMPLE AREA LOCATION - Section 3, Township 14 North, Range 1 East, MPM,
About three (3) miles west of Turtle Butte.

SUGGESTED POLICIES -

1. Grazing and other agricultural activities should be utilized to the fullest.

RECOMMENDED ACTIONS -

1. If any development takes place, extensive on-site evaluations should be made. Criteria weighed should be;
 - a. Soils limitations
 - b. Ground and surface water: depth to groundwater
 - c. Geologic hazards
 - d. Slope
 - e. Adverse effect on wildlife habitat
 - f. Visual impact
 - g. Effects on recreational uses and access
2. Land-use controls should be based on the aforementioned criteria.
3. Agricultural uses should be given top priority.

IGNEOUS UPLANDS - LANDSCAPE UNIT

DEFINITION - Anything above the 5000 foot contour.

EXTENT AND DESCRIPTION - The uplands landscape unit occupies the southwestern corner of the Planning District. The volcanic formations account for the areas ruggedness and isolation. Use of this land is for marginal livestock grazing. Forest can be found throughout the area.

VEGETATION - Vegetation is light in the uplands due to poor soils. Timber consists of various types of spruce, fir and pine. Quaking Aspen is often found in gulches of the area. Grass in the uplands is often bunchgrass, with juniper and rabbit brush also inhabiting the area.

WILDLIFE - Big game animals found in the upland landscape unit include: elk, mule deer, whitetail deer, Rocky Mountain Goat, and Rocky Mountain Sheep. Game birds are mainly grouse. Eagles and osprey can be found in those uplands near rivers and streams.

GEOLOGIC HAZARDS - Hazards in this unit consist mainly of landslides and rock falls. The steep terrain combined with the igneous formations make it particularly hazardous. Earthquakes, although no faults are known in the area, could work as a triggering action for slides and falls.

CHARACTERISTICS -

GEOLOGY - The geology of this unit is basically composed of Adel Mountain Volcanics which are potash-rich volcanics and related intrusive rocks. They are of the Late Cretaceous (60 million years ago) or early Quaternary Period (20 million years ago) and intrude into the Cretaceous strata of the area. The formations are highly associated with the colluvium alluvium of the foothills landscape unit. The igneous rocks are believed to have been intruded before the main diastrophism of the Disturbed Belt.

SOILS - Mucet-Sangrey-Rock outcrop association soils are found in the mountainous areas along the west side of the county and in the Mission Mountains. The soils are generally less than 20 inches deep to igneous bedrock. Mucet soils are dark colored with gravelly loam surface layers over gravelly loam subsoils. Depth to igneous bedrock is between 10 and 20 inches. The steepness and depth to bedrock puts severe limits on this soil for most uses.

Sangrey soils are generally of thick black loam and gravelly loam surface layers and brown loam subsoils. These generally lie under the Mucet soils and are frequently dissected by deep drainageways. Limitation are not too severe for this soil.

Rock outcrops are of barren igneous rock in irregularly shaped areas. The Mucet-Sangrey-Rock outcrop association is used primarily for grazing and woodland. Nonirrigated grain and hay crops are grown on Sangrey and other deep soils on the terraces and fans.

The Hanson-Dumco-Faines association is present but is of minor importance in the igneous uplands. This association is discussed in the sedimentary foothills landscape unit.

SLOPE - Slopes vary up to 75%.

EXAMPLE AREA LOCATION - The Sawteeth in Section 32, Township 16 North, Range 1 West.

SUGGESTED POLICIES -

1. Investigations for slope suitability, conducted by a professional engineer or geologist, should precede any construction.
2. Planning standards should reflect the preferred use of this landscape unit as being open space and wildlife habitat with grazing the primary land use.

RECOMMENDED ACTIONS -

1. The county should permit residential development in the uplands landscape unit only where there is little adverse impact on the environment.
2. An environmental impact statement should be required prior to any development.
3. All development should reflect the lay of the land.
4. Extensive on-site evaluations should be made prior to any development.

BUTTES - LANDSCAPE UNIT

DEFINITION - The conspicuous hills with steep or precipitous sides which are visible north and west of Cascade. Differential erosion has caused the more resistant capped buttes to stand above the surrounding landscape.

EXTENT AND DESCRIPTION - Buttes are one of the more unusual landscape features of the area. They are found extensively northwest of the Missouri River in the foothills and benches landscape units. Elevation is usually 300 feet to 1000 feet higher than the surrounding land. Notable buttes are Cascade Butte, Lionhead Butte, Haystack Butte and Johnson Butte.

VEGETATION - Vegetation is scarce due to the absence of soils in the landscape unit. Native grasses, cactus and sage inhabit the tops of these landforms.

WILDLIFE - Little wildlife is known to exist on the buttes. Eagles can be found on some of the inaccessible buttes but little habitation is known otherwise.

GEOLOGIC HAZARDS - Hazardous conditions in the buttes landscape unit are mainly landslides caused by breakdown of the formation from natural processes. Square Butte in the Sun River Valley Planning District has a large landslide on its south face. This particular slide is believed to have been caused by moisture in an extremely wet year which provided lubrication.

CHARACTERISTICS -

GEOLOGY - Within the Cascade Planning District a number of unusual buttes exist. The geology of these formations are intrusive igneous bodies called sills. They are formed by magma being forced between sedimentary beds causing a blister shaped formation. They are relatively flat in nature and are much more resistant to erosion than the surrounding landscape. This accounts for their prominence above the adjoining land. They are Tertiary (40 million years ago) developments as are all other igneous bodies in the area.

SOILS - Mucet-Sangrey-Rock outcrop soils are generally less than 20 inches deep to igneous bedrock. Mucet soils are dark colored, well drained, and free of lime. They have gravelly loam surface layers and gravelly loam subsoils underlain by igneous bedrock. Depth to bedrock and steepness limit the suitability of this soil for most uses.

Sangrey soils have thick black loam and gravelly loam surface layers and brown loam subsoils. These soils generally occur on fans and footslopes with slopes up to 30% and lie below Mucet soils. The slope of the soils is a minor limitation for some uses.

The rock outcrops are barren igneous rock in irregular shapes. The Mucet-Sangrey-Rock outcrop association is used mainly for grazing and woodland. Nonirrigated hay and grain crops are grown on Sangrey and other deep soils found on terraces and fans.

SLOPE -Relatively gently sloping tops with steeply sloping sides up to 90° near the crest.


EXAMPLE AREA LOCATION - Cascade Butte in Township 18 North, Range 1 West
Birdtail Butte in Section 2, Township 18 North, Range 3 West
Haystack Butte in Section 1, Township 18 North, Range 3 West

SUGGESTED POLICIES -

1. Primary Land use should be left as grazing.
2. Development should not be allowed.
3. Efforts should be made to protect the natural beauty of this landscape unit.

RECOMMENDED ACTIONS -

1. Land-use controls should be developed which reflect the environment and aesthetics of the buttes.
2. Buttes should remain undeveloped and incentives or sanctions should be designed through the planning process to maintain the buttes in an undeveloped state, possibly through public ownership.



LANDSCAPE UNITS

SECTION II

CRITICAL HAZARD AREAS

ENVIRONMENTAL PLAN FOR THE CASCADE PLANNING DISTRICT

SECTION II - CRITICAL HAZARD AREAS

The intent of the hazard areas section is to identify and record areas of hazardous nature in Cascade County, and to establish a procedure whereby proposed uses of these areas can be examined by citizens and county planners in order to protect life and property.

This section of the environmental plan describes areas which could be or are known to be susceptible to mass movement, flooding or earthquake. Since these pose risks for the construction of buildings, utilities, roads and the safety of persons living in the area, policies and performance standards are suggested for the various areas involved.

Some or all of the following information is provided for each hazard:

- a. General description
- b. Classification of the hazard areas
- c. Suggested policy and recommended actions
- d. Performance standards
- e. Possible procedure for identifying hazard areas.

The recommended policies and performance standards should serve until adequate floodway management and delineation is accomplished. At this time there is little available information on hazardous areas in Cascade County. Areas that are prone to flooding are generally known but have yet to be officially delineated. Areas that are susceptible to mass gravity movements have not been mapped or documented.

FLOOD HAZARD AREAS

GENERAL DESCRIPTION - Flood insurance regulations differentiate two (2) types of flooding, the "floodway" and the "floodway fringe." The "floodway" is the flood channel that carries the fast moving flood waters and is the area where flood damage will be the greatest. The "floodway fringe" is the area inundated by rising waters. Damage in the floodway is caused by fast flowing currents, while damage in the floodway fringe is caused by water damage and deposition of mud. The Missouri River and some of the smaller creeks have been known to flood.

DEFINITIONS -

FLOOD - The temporary rising and overflowing of a body of water onto normally dry land.

FLOODPLAIN - A plain built up by stream deposition which may be submerged periodically by floodwaters.

FLOODWAY - The normal stream channel and that adjoining area of the floodplain needed to convey moving waters at flood stage.

FLOODWAY FRINGE - The area of the floodplain lying outside the floodway, but subject to periodic inundation from flooding.

CLASSIFICATION OF FLOOD AREAS - There have been no designations prepared to classify lands in flood prone areas at this time. These designations are presently being considered: flood areas, floodway, and floodway fringe.

SUGGESTED POLICIES -

1. Housing and industrial developments should be discouraged in areas identified as flood prone.
2. Problems with flood prone areas should be handled as land-use considerations, not as engineering problems.
3. Hospitals, nursing homes, and schools should not be allowed in flood prone areas.
4. Through the use of performance standards and floodplain regulations, agriculture, horticulture, and other open space uses should be encouraged.
5. Construction of additional flood control measures should be discouraged. (Land-use problems rather than engineering problems)

RECOMMENDED ACTIONS -

1. All developments and subdividing of land should be discouraged, until floodplain designations are finalized.

2. A floodplain information and mangement plan should be developed by the county to inform the citizens of the hazards in these areas.

SUGGESTED PERFORMANCE STANDARDS -

1. The construction of access roads, embankments and fill should not impede the natural surface drainage pattern unless the express purpose of the contruction is to impound water.

ICE JAMS

GENERAL DESCRIPTION - Ice jams are formed when ice builds up and packs

together, forming a "dam" that obstructs river flow. Flooding may result. Slab ice moving onto the alluvial lowlands may push against obstacles in its path, exerting enough pressure to move a house off its foundation. Most damage is caused by water, rather than the ice itself. Ice jams are unpredictable, both in place occurrence and conditions necessary for formation.

Most ice jams in the Cascade Planning District occur on the Missouri River. Areas near Seibold and Willington tracts in the southwestern part of the county have been known to flood on several occasions when water was trapped behind an ice jam further down river. Information on ice jams is generally scarce unless damage results.

DEFINITIONS -

ICE JAMS - A buildup of tightly packed ice that obstructs normal stream flow.

CLASSIFICATION OF ICE JAM AREAS - There have been no designations prepared to classify lands in areas prone to ice jams. Designations should be made on the basis of possibility rather than probability of occurrence because of the extremely unpredictable nature of ice jams.

SUGGESTED POLICIES -

1. Housing and industrial developments should be discouraged in areas of known ice jam occurrence.
2. Hospitals, nursing homes and schools should be prohibited in areas of known ice jam occurrence.

RECOMMENDED ACTIONS -

1. All occurrences of ice jams, whether damaging or not, should be reported to the Cascade County Planning Board and recorded.
2. Purchasers of land in these areas should be informed of the potential hazard involved.

MASS MOVEMENTS

GENERAL DESCRIPTION - Mass movement is the downslope movement of materials in response to gravity. Because of the nature of some areas of the county, there exists the potential for the mass movement of earth and rock. Contributing factors to the problem of hazardous mass movements include: slope, geology, soils, vegetation, rainfall (intensity and duration), man-made and natural disturbances and the variable relations and combinations of these factors. Magnitude and material content of mass movements vary with location and can be characterized by rock fall, soil creep, earth flow, slumping bedding plane failure, debris slide or flow, or mudflow.

Hazardous mass movement areas can be found throughout the Cascade Planning District. Slumping and soil creep are evident in all landscape units and are the primary mass movement hazards in alluvial lowlands and benches and terraces. Rapid mass movement such as rock falls and debris slides are more common in uplands, foothills and buttes, but can sometimes be found on the steeper slopes of the lower landscape units.

DEFINITIONS -

BEDDING PLANE - The junction plane between two (2) strata or layers of rock.

BEDDING PLANE FAILURE - Sliding along planes of weakness where bedding planes are parallel to the dip (slope). When only rock is involved the term "rock glide" may be used.

The phenomenon of bedding plane failure is common where shale is overlain by sandstone and is aggravated where lubricated by water.

DEBRIS SLIDE - Rapid downslope movement of loose rock or soil on steep slopes.

EARTH FLOW - The movement of a mass of soil or of rock and soil. A scarp usually develops on the uphill side of the flow and soil, rocks, and vegetation are mixed together. Shallow sag ponds and irregular drainage patterns are common.

MUDFLOW - The rapid, relatively fluid flow of soil and rock under conditions of excessive saturation.

ROCK FALL - The process by which rock becomes detached from a nearly vertical slope resulting usually in talus slope.

SLUMPING - The process by which a more or less continuous mass of earth slips downhill along a curved basal slip plane in such a way that the mass of earth rotates backwards, toward the slope. The top surface of a slump frequently has a backward slope with a scarp on the uphill side and a toe on the downhill side. This is usually a relatively rapid movement.

SOIL CREEP - The very slow downslope movement of soil and weathered residuum. Frequently, this type of movement can be identified by the bowing of trees or the tilting of fences and utility poles.

For the most part, mass movement areas are easy to identify, as very unique landscape results. However, where the movement is very slow or is located a distance from the sight, identification becomes much more difficult. In addition, movement may be initiated or accelerated by man's activities. Construction activities and facilities frequently change the surface drainage, the infiltration of water, and the load, thereby increasing the risk of movement. Excavation can expose a bedding plane that is susceptible to movement or may oversteepen a slope. Cutting the toe off an old slide can make it unstable and reinitiate movement.

CLASSIFICATION OF MASS MOVEMENT AREAS -

1. Critical mass movement areas - Areas that currently display mass movement.
2. Potential mass movement areas - Areas which have displayed mass gravity movement in the past and have the potential to become active. These areas may be suitable for limited development but extensive on-site evaluations should be performed prior to any action.

SUGGESTED POLICIES -

1. Hospitals, schools, nursing homes and community centers should be prohibited in areas displaying mass movements.
2. Purchasers of land in areas designated as subject to mass movement should be informed as to the potential hazards inherent to those areas.
3. All construction activities should be discouraged in areas presently expressing mass movement or a high potential for such.
4. Potential development of areas of landslide topography should be scrutinized to assess its affect on the land; those uses which may have a detrimental affect should be discouraged.

5. Extensive on-site investigations should be made by a competent expert to determine where a potential hazard exists.

RECOMMENDED ACTIONS -

1. An effort should be made to accumulate all available geologic information.
2. Performance standards should reflect the environmental constraints of mass movement areas.
3. Efforts should be made to inform potential buyers of the inherent hazard areas.

SUGGESTED PERFORMANCE STANDARDS -

1. No new structures should be permitted in areas identified as critical mass movement areas.
2. No cut and fill operations should be permitted.
3. In areas of potential mass movements, a site inspection should be carried out by a soils engineer or geologist.

PROCEDURE FOR IDENTIFYING MASS MOVEMENT -

1. Existing reports
2. Field investigations by qualified professionals
3. Montana State Highway Department - Engineering Division

EARTHQUAKES

Earthquakes present a potential hazard in the Cascade Planning District. Fault lines generally follow a northwest - southeast directional trend and are primarily located in the southern part of the district.

Earthquakes have been recorded as close as Craig, Montana in Lewis and Clark County but none are known to have been centered within the county. The Carter Ranch Fault is one of the largest faults in the area. It is located in Township 16 North, Range 2 East and runs for a total of 9½ miles. Displacement ranges up to 700 feet at several points along the faultline.

SUGGESTED POLICIES -


1. Plans for development in the proximity of fault lines should acknowledge the presence of this hazard, and all potential buyers should be made aware of its existence.
2. The presence of faults in the area should constitute an additional reason for restricting development in areas of landslide topography.

RECOMMENDED ACTIONS -

1. Encourage site examination by a qualified engineer or geologist in areas of known faults.
2. Discourage uses such as schools, hospitals, churches, and large residential developments in the proximity of fault lines.



CONTOUR INTERVAL: 1000 ft



TOPOGRAPHY

SECTION III

OPEN SPACE

ENVIRONMENTAL PLAN FOR THE CASCADE PLANNING DISTRICT

SECTION III - PUBLIC OPEN SPACE

PART I - EXPLANATION

Open space in all of its many forms has increased in importance to such a degree that it has gained recognition as one of the major resource management functions. The contributions of open space to human well-being are becoming widely recognized, and public awareness of dwindling open space resources is growing. As the population and attendant urbanization proliferates, so privacy, solitude, and the availability of uncrowded recreational areas become more elusive. In addition, public attitudes are becoming more sophisticated in accepting the importance of the various ecological functions of certain areas when kept in their natural state. The new attitudes extend to the recognition that some areas are inherently unsuitable for development because of geologic or climatic instability, and that these areas should be preserved in open space because of the potential hazards to human life or property.

Because of their financial dependence on property tax, local governments have typically favored development over the preservation of land as open space, but the belief that tax rolls are benefitted more by development is currently being subjected to critical reexamination as a result of recent studies. A number of these studies concluded that to allow land in certain areas to be developed would prove more costly to the public than to acquire the land for public open space. Furthermore, the economic arguments used formerly to promote development or to oppose open space proposals are no longer being accepted at face value by the public or local governments.

In the environmental plan, public lands which possess significant existing or potential value for public open space purposes are identified and recommended for preservation for future open space uses. In arriving at the individual sites being proposed in the open space plan, such factors as access, usability, vegetation cover, historical value, and recreation were considered. It must be remembered that sites shown in the plan are not the only possible sites and may warrant adjustment or deletion as time and other factors may dictate.

It is our objective in this section to identify and classify existing and potential open space areas in the Cascade study area, along with suggested goals, policies, and recommended actions for the Citizen's Advisory Committee and the Board of County Commissioners to consider.

PART II - DEFINING OPEN SPACE

The term "open space" is an abstraction which eludes simple definition. Open space may be defined legally, as for legislative purposes; or functionally, defining a use for space.

A. LEGAL DEFINITIONS OF OPEN SPACE:

Some ordinances and laws have defined open space for a number of purposes. Some definitions indicate types of land for which local governments can receive grants. Others indicate the types of land to be considered in the preparation of local open space plans.

The federal definition of open space is contained in the open space provisions of the Housing Act of 1961: "The term 'open space land' means any undeveloped or predominately undeveloped land in an urban area which has value for (a) park and recreation purposes, (b) conservation of land and other natural resources, or (c) historic or scenic purposes."

Montana's model subdivision regulations define open space as "A land or water area devoid of buildings and other physical structures except where accessory to the provisions of recreation."

B. FUNCTIONAL DEFINITIONS OF OPEN SPACE:

Open space is often identified as land which does or can serve some use as open space. For example, Cascade County's Subdivision Regulations contain the following:

"Common Open Space: A parcel or parcels of land, or an area of water, or combination of land and water within a planned unit development designated or intended for the use or enjoyment of residents within the development."

In this context the definition consists of the function which open space serves. In the Cascade County Open Space Program, the following open space functions will be used to organize both the inventory of the county's open space resources and the set of findings and policies regarding open space:

1. Open space for recreation; parks, park lands, trails and water areas.
2. Open space for the protection of scenic resources; scenic backdrops, scenic roads and highways, and scenic water areas.
3. Open space for the protection of wildlife and vegetation.
4. Open space for the protection of historic, archeological and scientific resources; historic landmarks and archeological sites.

In all likelihood, not all of the above open space designations will be used in any one (1) planning district.

PART III - NEED FOR OPEN SPACE:

Residents in Cascade County are fortunate in that they are within easy travel distance of many recreational opportunities. Glacier National Park; Holter, Hauser, and Canyon Ferry Reservoirs; the Missouri, Sun, and Smith Rivers; Showdown Mountain Ski Area; the Bob Marshall and Gates of the Mountains Wilderness areas; and a rather large portion of the Lewis and Clark National Forest are right on the doorstep. According to the State of Montana Fish and Game Commission demand (in terms of occasions per weekend) for Region 5 was 185,300 activity occasions. Region 5 includes Glacier, Toole, Pondera, Chouteau, and Cascade Counties. The increase demand projections for selected activities by 1990 are:

Camping - 44%	Hiking - 46%
Picnicking - 23%	Driving - 22%
Boating - 48%	Sight-seeing - 29%
Swimming - 38%	Walking - 28%
Fishing - 18%	

At present, excluding Glacier National Park, the following facilities are in Region 5:

- 1) 57 picnic units
- 2) 141 camping units
- 3) 1 swimming unit
- 4) 16 fishing access sites
- 5) 4 boat ramps

With population growth and increasing amounts of leisure time, a growing demand for open space and recreation activities seems to indicate a need for additional facilities. There is a lack of facilities close to the smaller communities. Facilities suitable for day use activities in areas of a natural environment are deficient, especially near Great Falls.

This demand can only be met by designating and setting aside lands, both public and private, to be used and developed into future recreational and open space sites. Designation and reservation of land should begin as soon as possible as the demand for residential as well as other types of development is likely to occur simultaneously.

Areas which should receive special attention:

1. Fishing access and camping sites along the Missouri River.
2. St. Peter's Mission - ten (10) miles west of Cascade. Mission for educating the Blackfeet Indians.

3. Mt. Palace - about ten (10) miles southwest of Cascade.
Unusual geologic formation.
4. Buttes - throughout the Cascade Planning District. Unusual
geologic formations.

Another important area of potential open space includes what is locally known as Tintingers Slough located several miles south of Cascade. The area is a wooded lowland and oxbow lake formed from a former meander of the Missouri River. Light flooding is common in the area making it impractical for residential type developments. The land is presently used as pastureland with some small haylands. Interstate 15 bisects the area making it an excellent location for combination rest-stop and a county park. Presently this land (as are the buttes) is under private ownership, and acquisition, if possible, would be through a mutually acceptable transaction.

U.S. Highway 91 from Cascade to Helena could be prospective public open space project. Designation as a scenic highway would protect it from defacement and would preserve much of the scenic landscape of the area. Bike-ways, information signing and scenic viewpoints along the highway could be added attractions, as the road is now used primarily for recreation and fishing access. Funds for financing this type of project are available through both Federal and State agencies and a joint project could possibly be worked out with Lewis and Clark County.

A. OPEN SPACE FOR RECREATION:

FINDINGS -

1. Because they are held in public ownership, parks are a permanent form of open space that can be counted on to endure from generation to generation.
2. The private sector can play an important role in providing recreational open space areas.
3. Open space recreational developments, if not properly designed, may have an adverse effect on natural resources and the environment in general.
4. Projects of special districts (e.g. flood control districts) and utility companies can sometimes be designed and constructed to include trails along the right-of-way of the project.
5. Recreation activities on rivers and streams, if not properly regulated, can be harmful to important wildlife species and wildlife habitat areas.

SUGGESTED GOALS -

1. Develop and maintain recreational opportunity for groups of all ages.

SUGGESTED POLICIES -

1. Proposed recreational developments that may have an impact on the environment shall be carefully designed and reviewed and environmental impact assessments should be made.
2. Use of off-road motorized vehicles shall be generally prohibited in open space areas.
3. Additional public access and appropriate recreational facilities shall be provided along rivers and streams.
4. Provide recreational facilities capable of promoting the health, safety, morals and general welfare of the public.
5. Certain lands in public ownership shall be set aside for future use and development as open space, as there is a need for them because of population and industrial growth.
6. Land exchanges between public agencies and private owners shall be encouraged for the acquisition of open space when applicable.
7. Provide a diversified year-round recreation plan and program to serve the residents of Cascade County.

8. Recognize the potential of the region as a tourist recreation area and provide for a balanced development of tourist oriented facilities which will enhance and stabilize the region's economic base.
9. Water-dependent recreational development shall be considered a priority use.
10. Water-dependent recreational development providing public access shall be given preference over similar development which limits public access or use.
11. Recreational development and use shall not exceed the carrying capacity of the land.
12. Adequate public access to river and stream resources shall be provided.
13. Public access roads and trails shall be provided to areas currently in public ownership.
14. Use existing public land for open space before acquiring additional sites.

RECOMMENDED ACTIONS -

1. Local and state governments should work actively to provide parks and other recreational open space.
2. Provisions should be made to encourage the private sector to enter the field of outdoor recreation.
3. Programs for park acquisitions and development should include lands for trails. Trails should be built as soon as possible after acquisition, using volunteer help for construction and maintenance wherever possible.
4. The State Division of Highways should be urged to include right-of-ways for bicycle trails where appropriate along state highways.
5. State, federal, county, and city park and recreation agencies should be encouraged to coordinate their efforts.
6. Additional study should be done to determine a stable and continuing source of funds for park acquisition and development of appropriate scale to adequately support anticipated needs for recreation and park service.
7. A map listing fishing access sites should be developed cooperatively by the County Planning staff and the Montana Fish and Game Commission.

B. OPEN SPACE FOR THE PROTECTION OF SCENIC RESOURCES:

FINDINGS -

1. Cascade County is rich in number of roads and highways traversing scenic rural areas. Leisurely travel on these roads is a valuable recreation and open space experience. Some roads should be given protection against defacement by scenic highway zoning.
2. Increasing travel on scenic roads may greatly diminish their value.
3. The scenic value of water areas can be greatly diminished if development is allowed to encroach upon them.
4. Open spaces within the urban area are valuable for the scenic enjoyment of urban dwellers.

SUGGESTED GOALS -

1. Encourage preservation of the scenic beauty and natural environment.

SUGGESTED POLICIES -

1. River access shall be maintained for public use.
2. Great care shall be taken to protect the character of scenic roads.
3. Areas adjacent to bodies of water shall be protected for their value as scenic open space.
4. Maximum pedestrian access shall be provided for scenic enjoyment of the open water; sloughs, marshes, and rivers - except for natural areas protected for their scientific and educational value.
5. All county lands to be sold at public auction into private ownership should be reviewed by the Cascade County Planning Board and Citizen's Advisory committees for potential use and public open space lands before said sale.
6. Use of land for open space shall be considered a priority use of the study area.
7. Wherever practicable, open space shall be enhanced and protected to insure retention of the existing local character.
8. Scenic vistas of the study area shall be designated and protected.

RECOMMENDED ACTIONS -

1. There should be a study of scenic roads to determine their suitability for recreation trails, turnouts, and parking areas.
2. Where appropriate, scenic easements should be acquired from private landowners.

3. The county should provide standards and programs for maintenance of open space and protection of scenic areas through development criteria.

C. OPEN SPACE FOR THE PROTECTION OF WILDLIFE AND VEGETATION:

FINDINGS -

1. Open space areas are important to wildlife in providing them with the food, resting, and nesting areas needed for their survival.
2. Each species of wildlife is adapted to and dependent upon a particular set or range of habitat conditions for its survival. Some species can survive urban or suburban areas, but others can survive only if their habitat is preserved in its natural state, largely undisturbed by man. Even those which can survive in urban areas may be able to do so only so long as there remains within the urban areas predominately natural areas, such as wooded stream sides, where they can find refuge and their other survival needs.
3. Several species of wildlife in Cascade County are included on the U.S. Department of Interior's list of rare and endangered species. If their habitat areas are not carefully protected, these species may become extinct.
4. The opportunity to observe and study wildlife in their natural habitat is a source of leisure time enjoyment to many residents of the county. Human intrusion into wildlife areas, however, can have adverse impact upon wildlife and their habitats.
5. Vegetation performs a number of important open space functions, including:
 - (a) Providing scenic beauty
 - (b) Serving as habitat for wildlife
 - (c) Preventing soil erosion
 - (d) Maintaining air quality
 - (e) Controlling water runoff in watershed areas.
6. Certain types of recreational activity, such as the use of off-road vehicles, can adversely affect vegetation in open space.

7. Natural vegetation is an irreplaceable education resource as it allows study of features and conditions of natural environmental balances.

RECOMMENDED ACTIONS -

1. Retention of vegetation in both urban and nonurban areas should be a major objective of local open space and environmental policies.

D. OPEN SPACE FOR THE PROTECTION OF HISTORIC, ARCHEOLOGICAL AND SCIENTIFIC RESOURCES:

FINDINGS -

1. Historic landmarks and archeological sites may be lost if they are not protected from urban development.

SUGGESTED GOALS -

1. Preserve, develop, and maintain historical, educational and natural sites.
2. The overall goal in developing policies regarding historical and archeological resources is to preserve as a living part of the life and development of local area, region, state, and nation the rich heritage of our past that might otherwise be lost due to unnecessary and unwise development.

SUGGESTED POLICIES -

1. Both public and private efforts shall be directed to preserve historical landmarks which have open space potential.
2. Areas and sites of archeological significance shall be protected from development.
3. The public interest in areas of unique historical, scenic or archeological significance shall be recognized in considering development in these areas.

RECOMMENDED ACTIONS -

1. Historical and archeological resources should be inventoried and protected.
2. The county should evaluate the impact of development upon historical and archeological resources prior to approval or disapproval of the project.

PART IV - IMPLEMENTATION

The following implementaion techiques offer possible ways of acquiring future open space in Cascade County assuring the maintenance of our rural quality of life.

1. Property Donation. Property donation is a way for an individual to give the county a piece of property for whatever reason he wishes. Should this property be suitable for open space use, the county could keep it in reserve for future use. This, of course, is a voluntary method of acquiring open space land.

2. Property Exchange. When the county wants a piece of property which is owned by a private individual or corporation, the county can offer to exchange part of the land now in public ownership for this privately owned parcel.

3. Conservation Easement. A conservation easement is a piece of land privately owned on which certain rights have been sold or donated to the city, state, or county by the owner. The rights sold or donated would be tied to the land and would forbid certain types of building and development. This would be a voluntary agreement between the public body and the private individual.

4. Sale-back and Lease-back. In this method of acquiring land for open space, the county buys the land outright and places property restrictions and covenants on the land. The land is then sold or leased back to the private sector.

5. Cluster Development. This system allows the county and land developer to negotiate an agreement whereby the developer may increase the density beyond the stated maximums in the county zoning ordinance on one part of his property and leave another part free from development as publicly dedicated open space.

6. Club or Group Involvement. Civic, service and garden clubs, Boy Scouts and military construction units, can make important voluntary contributions to public recreation areas. When a park development plan is available, such groups can concentrate on projects which interest them and gradually bring about improvements to facilities even when public funds are not available.

7. Private Recreation Facilities. When private facilities answer an obvious public need and are properly managed, the county should encourage and supplement, rather than duplicate them.

8. Easement, Lease, Development Right Purchase. The purchase of development rights is one device for keeping land in its existing condition. The owner may continue to use the land as he has in the past but may not develop it for some new, more intensive use.

9. Use of Tax Foreclosed Land. In undeveloped areas where recreation areas are needed, the possibility of utilizing tax foreclosed lands should be investigated. In some cases, lands owned but not used or needed in the future by some other governmental agency may also be put to temporary or permanent recreational use.

10. Continuing Levy. An alternative to a special election for open space acquisition is to obtain voter approval for continuing levy to be applied to open space acquisition. This assurance of a regular revenue permits a broader development program to be planned and accomplished.

11. Federal Financial Assistance. A number of federal programs have been established which might prove helpful to the county in financing open space acquisition and improvements.

12. Fee Simple Acquisition. Acquiring land outright for open space by the use of federal, state, or local funds, is another method for acquiring open space lands.

SECTION IV

PRIORITY RESOURCES

ENVIRONMENTAL PLAN FOR THE CASCADE PLANNING DISTRICT

SECTION IV - PRIORITY RESOURCES

The purpose of this section of the plan is to identify priority resource areas within the district and to provide a sound policy foundation for conservation policies. Priority resources may be defined as lands containing natural resources of either existing or potential economic value. The criterion for identifying priority resources is any resource which serves an important economic function. Included as priority resources in the Cascade Planning District area: agricultural lands; fisheries; mineral and mineral aggregates; and water resources.

AGRICULTURAL LANDS

Agriculture is the primary land use in the Cascade Planning District. The area contains some excellent irrigated lowlands, dryland grain farms and pasturage for livestock. Sprinkler irrigation has recently been introduced on some of the dryland farms bringing higher production rates to the area.

Certain parts of the district are presently being transformed from agricultural to suburban and rural residential areas. (The Missouri River lowlands southwest of Cascade is a notable example.) Lands in transition are probably not the most agriculturally productive, but as encroachment upon these areas continues, adjacent or nearby agriculturally productive lands are pressured by the higher values of residential and industrial uses. Eventually they succumb to further development resulting in loss of agricultural land and continuation of the transformation process.

SUGGESTED POLICIES -

1. Rather than allow indiscriminate leap-frog subdivision and development, residential subdivisions should be developed in clusters near or adjacent to present urban or residential areas.
2. Efforts should be made to preserve lands now under irrigation; including those lands which have a high potential for being irrigated at some future date.
3. Floodplains should be left for agricultural and open space uses.
4. The aesthetic value of agriculture should be recognized and agricultural uses should be encouraged along scenic corridors.
5. Local land-use planning should encourage the continuance of agriculture as a functioning part of the local economy.
6. Agricultural activities and practices which are compatible with the physical suitability and landscape of the planning district should be encouraged.

RECOMMENDED ACTIONS -

1. Local governments and others should discourage conversion of agricultural land to urban uses unless it is an orderly and necessary extension of an existing land use.

FISHERIES

Although not directly a land resource, sport fishing is directly or at least partially affected by activities on the land adjacent to water bodies. Commercial fishing is not and probably will not be an important in the near future in Cascade County. The Missouri River is classified as a cold water sport fishing stream.

SUGGESTED POLICIES -

1. No waste shall be discharged into any waters without first receiving the treatment necessary to protect the fisheries resources.
2. Buffer strips should be encouraged to keep debris out of active flowing streams.
3. Rechanneling of streams should not be allowed.
4. Adequate flows shall be maintained in streams in order to provide habitat for fish.

RECOMMENDED ACTIONS -

1. The county should establish an environmental assessment procedure requiring extensive on-site investigations before any development takes place within 100 feet of the river bank. These assessments should be made public and reviewed by qualified technical personnel to judge their accuracy.
2. Minimum stream flows should be established to protect and provide habitat for fish.

MINERALS AND MINERAL AGGREGATES

Sand, gravel, and crushed rock are important factors in the development of any community. These mineral aggregates are used in the making of portland cement concrete, asphaltic concrete, foundations for roads as well as select fill and are used extensively in concrete for large structures such as bridges and multi-story buildings. There are several areas in the planning district where sand and gravel are presently or have been mined with other areas having the potential to produce building materials.

Other potential minerals found in the area are sandstone, alluvial clay, coal, oil and gas. Sandstone in small amounts is used for foundations, well linings, decorative stone, and riprap for facing on small stock dams. Some of the alluvial clay in the Missouri Valley has ceramic properties although it is not present in sufficient quantity to be of economic importance. Coal found in the western part of the planning district is in seams greater than fourteen (14) inches, which may or may not be economical to mine in the future. Hound Creek near the Smith River is a former coal producing area. There are several coal leases in the southern part of the planning district. There is a potential for oil and gas production in the southeastern part of the planning district. Also identified are areas presently under mineral lease; areas which the federal government owns all mineral rights and privately and federally owned land with partial federal ownership.

SUGGESTED POLICIES -

1. Because mineral resources are in short supply, all known areas of economic potential should be protected from any type of development that would conflict with extraction activities.
2. Require reclamation of areas after mining is completed.
3. Require an Environmental Impact Statement.

RECOMMENDED ACTIONS -

1. The county should identify areas having minerals with economic potential.
2. Exploration of available and potential mineral resources should continue and extraction and development encouraged if it is determined economically, socially, and environmentally acceptable.

WATER RESOURCES

The Missouri River flows through the Cascade Planning District from southwest to northeast and supplies most of the water needs of the area. Exact acreage of land under irrigation in this area is not available but several small projects have been developed in the Chestnut Valley and other lowland areas. A few projects have been abandoned such as an irrigation system on the Pistoria Tracts at the confluence of the Missouri and Hardy Creek. Hound Creek in the Smith River drainage has an irrigation project serving a total of 2,214 acres. Another irrigation project can be found on Little Muddy Creek and its tributary Rock Gap Creek. Over 1,500 acres are served by this project.

Many small creeks are diverted into ditch networks to utilize natural run-off. This type of irrigation accounts for much of the irrigation in the upper Chestnut Valley.

Excellent groundwater sources are available throughout the district. Notable aquifers include the Kootenai Formation which is one of the best aquifers in Montana. Little development of this aquifer has taken place in the Cascade area because it is generally beyond economic drilling depth. The Blackleaf and Telegraph Creek Formations also contain very good aquifers, several of which are suitable for irrigation purposes.

SUGGESTED POLICIES

1. No appropriations of water except for human consumption, live-stock consumption and water legally released from storage shall be granted by any state agency when the established minimum perennial stream flow is less than that to support aquatic life.
2. No out of basin appropriation of the waters of the Missouri River Basin shall be considered without the prior approval of, and following the public hearing in the subject basin by, the State Water Resources Board.
3. Rights to the use of water shall be granted only on the condition that any effluents or return flows shall not interfere with other beneficial uses of water.

RECOMMENDATIONS - If the streams in the study area are to remain an important resource, care must be taken to preserve this valuable commodity. As streams reach their low annual stream flow during the summer months, care must be taken to insure that the stream flow does not fall below the recommended minimum flow necessary to support aquatic life. State agencies should also regulate the issuance of water rights so as to insure that the total water rights of a stream do not exceed the minimum stream flow.

MISSOURI RIVER MAIN STEM
Missouri River near Ulm, MT

LOCATION - Lat 47°26'09", long 111°23'07", in NE $\frac{1}{4}$ NW $\frac{1}{4}$ NW $\frac{1}{4}$ sec. 5, T.19 N., R.3 E.,

Cascade County, on left bank 5.6 mi (9.0 km) east of Ulm and 9.1 mi

(14.6 km) downstream from Smith River.

PERIOD OF RECORD - August 1957 to 1974.

AVERAGE DISCHARGE - 17 years, 6,564 ft³/s* (185.9 m³/s), 4,756,000 acre-ft/yr (5.86 km³/yr).

EXTREMES - 1974: Maximum discharge, 20,000 ft³/s (566 m³/s) June 22, gage height,

12.09 ft (3.685 m); minimum daily, 3,220 ft³/s (90.6 m³/s) Oct. 1.

Discharge, in cubic feet per second, water year October 1973 to September 1974.

	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP
TOTAL	132,080	168,430	178,160	188,050	172,860	210,00	229,020	283,350	419,910	248,800	182,620	156,600
MEAN	4,261	5,614	5,747	6,066	6,174	6,490	7,634	9,140	14,000	8,026	5,891	5,220
MAX	4,990	6,030	6,000	6,600	6,660	7,400	10,300	12,600	19,900	17,600	7,140	6,730
MIN	3,220	5,080	5,450	5,100	5,800	5,890	6,060	7,560	9,860	5,510	4,330	4,260
AC-FT	262,000	334,100	353,400	373,000	342,900	399,100	454,300	562,000	832,900	493,500	362,200	310,600


CAL YR 1974 TOTAL 1,703,630 MEAN 4,667 MAX 6,950 MIN 2,240 AC-FT 3,379,000

WTR YR 1974 TOTAL 2,561,080 MEAN 7,017 MAX 19,900 MIN 3,220 AC-FT 5,080,000

NOTE - Stage-discharge relation affected by ice Jan. 1 to Feb.11.


* ft³/s = cubic feet per second.

MISSOURI RIVER MAIN STEM
Missouri River near Ulm, MT



MEAN DAILY FLOW PER MONTH
TOTAL MONTHLY FLOW

LEWIS AND CLARK CO



DRY LAND FARMING



HAY LAND



GRAZING



GRAVEL PIT



1 in = 4 mi

PRIORITY RESOURCES

APPENDIX A

DEFINITION OF LANDSCAPE UNITS

DEFINITION OF LANDSCAPE UNITS

RIVERS, STREAMS, LAKES AND RESERVOIRS

Lakes, reservoirs and rivers include water bodies either flowing or standing for all or most of the year. Included in this landscape unit are those lands immediately adjacent to water bodies which directly influence the physical, biological and chemical properties of the water.

ALLUVIAL LOWLANDS

The alluvial lowlands are plains occupying valley floors which result from the deposition of material by water. The major conspicuous feature of the landscape unit is the floodplain.

SEDIMENTARY TERRACES OR BENCHES

The relatively flat or gently sloping topographic surfaces which mark former valley floor levels. Stream downcutting has caused the terraces to stand higher than the present valley floor. Terraces are traditionally composed of alluvial deposits of sand, silt and gravel. Some of the necessary gravels, however, have either not been deposited by the stream or have been subsequently eroded. Therefore, the term "bench" may be more appropriate. There are some remnant terrace sand, silt and gravel deposits throughout the county, so the terms "terrace" and "bench" may be used interchangeably throughout the plan.

SEDIMENTARY FOOTHILLS

The foothills lie between the sedimentary terraces and the igneous uplands. The lower parameter is the 4000 foot contour line and the upper parameter, the break between foothills and uplands, is the 5000 foot contour line.

IGNEOUS FOOTHILLS

The igneous foothills occupy the area between the sedimentary terraces and the igneous uplands. The lower parameter is the same as the sedimentary foothills, the 4000 foot contour. The upper parameter corresponds with the 5000 foot contour.

SEDIMENTARY UPLANDS

The uplands landscape unit occupies much of the extreme southern part of the county with the 5000 foot contour as its lower parameter. Sedimentary uplands are those uplands composed of rock strata deposited by inland seas which once occupied the area.

IGNEOUS UPLANDS

Igneous upland are those areas underlain by igneous material and over 5000 feet in elevation.


METAMORPHIC UPLANDS

In Cascade County the metamorphic uplands landscape unit includes those areas geologically altered by volcanic or other activity. Metamorphic or metamorphism implies a pronounced change effected by heat and pressure resulting in a more compact, highly crystalline condition.

BUTTES

Buttes are the many conspicuous hills with steep or precipitous sides which are visible north of Cascade or south of Simms and Fort Shaw. Differential erosion has caused the more resistant capped buttes to stand taller than the surrounding landscape.

LANDSCAPE UNITS IN CROSSECTION



APPENDIX B

GEOLOGIC TIME SCALE

GEOLOGIC TIME SCALE

NAME		IMPORTANT EVENTS IN NORTHERN ROCKIES	
	Pleistocene Epoch	Ice ages and widespread volcanic activity	
- - - - - BEGAN ABOUT 3 MILLION YEARS AGO - - - - -			
	Tertiary Period	Valley-fill deposits laid down during time of desert climate. Mountain building, volcanic activity, intrusion of granites.	
- - - - - BEGAN ABOUT 60 MILLION YEARS AGO - - - - -			
MESOZOIC ERA	Cretaceous Period	Beginning of formation of Rocky Mountains	
	Jurassic Period	Volcanic activity, intrusion of granites	
	Triassic Period	Deposition of Mesozoic sedimentary rocks	
	- - - - - BEGAN ABOUT 225 MILLION YEARS AGO - - - - -		
PALEOZOIC ERA	Permian Period	Deposition of Paleozoic sedimentary rocks	
	Pennsylvanian Period	Formation of coal beds	
	Mississippian Period		
	Devonian Period		
	Silurian Period		
	Ordovician Period		
	Cambrian Period		
- - - - - BEGAN ABOUT 600 MILLION YEARS AGO - - - - -			
PRECAMBRIAN ERA		Deposition of Precambrian sedimentary rocks	
		Formation of Precambrian igneous and metamorphic basement rocks	
- - - - - BEGAN ABOUT 3000 MILLION (3 BILLION) YEARS AGO - - - - -			

APPENDIX C

ENDANGERED SPECIES

ENDANGERED SPECIES FOUND IN MONTANA

BLACK-FOOTED FERRET - may possibly be found in Cascade County but no research has yet been done.

PEREGRINE FALCON (Duck Hawk) - may be found in Cascade County but is very rare.

WHOOPING CRANE - none found in Cascade County.

NORTHERN ROCKY MOUNTAIN WOLF - none known in Cascade County.

The Black-footed Ferret is an extremely rare animal that eats prairie dogs. Its primary danger is that it may be poisoned when prairie dogs are poisoned.

Poisoning is also a threat to the Peregrine Falcon. Certain insecticides, such as DDT, concentrate as they move up through the food chain. The Peregrine Falcon is at the top of one such food chain where concentrations are very high.

APPENDIX D

ALTERNATIVE ENERGY

ALTERNATIVE ENERGY

The 1970's have introduced a new public awareness of the problems plaguing energy production in the United States. Shortages caused by excessive demands, political variables, and depletion of nonrenewable natural resources have shown that we are facing an uncertain energy future. Because planning is preparing for the future, alternative energy sources which are compatible to the environment and the public of Cascade County should be analyzed.

Present energy consumption within the county is mainly hydroelectric and natural gas. Various petroleum products are produced in local refineries and include transportation fuels and heating fuels. Wood is primarily used in the area for supplementary heating, but inefficient fireplace systems waste much of its potential.

Natural resources available within the county but undeveloped include coal, possible natural gas, sun and wind. The advantages and disadvantages of each resource are listed on the included energy resource cost chart. Energy resources unavailable in Cascade County but of importance to the nation are also listed.

Implementation of alternative energy methods generally falls into two (2) categories: private and public. Alternative private energy resources which can be developed on an individual basis include: solar heating, wind electrical generation, wood, and various others. This appendix will stress individual rather than community projects.

Because our conventional energy and fuel sources (oil, coal, natural gas and uranium) will not last forever, two (2) realizations must be made. The first is that we must slow down the consumption of such energy sources. The second is that we must find alternative sources to satisfy our tremendous energy appetite. In deciding which alternative energy sources to use we must choose those which would be the least costly and of the most benefit to America in the long run.

Following is a description of some of the nonrenewable and limited sources of energy available to the nation. Many of these are not located within Cascade County.

Nuclear Fission: Uranium is the fuel for these power plants and it is already in short supply. None is known to exist in Cascade County. Liquid Metal Fast Breeder Reactors are being developed which create more fuel than they can consume. Many think they are the hope for the future but they are still very controversial. Extremely high operating temperatures could cause explosions of poisonous material, and wastes generated by the system are difficult to dispose of.

Coal Gasification and Liquefaction: Coal reserves are a potential short term energy source in Cascade County. Commercial mining of this mineral was started as early as 1876 but has since been discontinued. Strippable deposits are located under much of the southern and eastern parts of the county. The environmental and social costs of developing this resource would be high.*

Oil Shale: Development of the oil shale in western states would impose heavy environmental costs in disruption of the land and in visual, air and water pollution. Not only is the waste material enormous but it would be of considerably greater volume than the original shale. Huge amounts of water are also needed. No oil shale is known to exist in Cascade County. This energy source is nonrenewable and is exhaustible although it would yield tremendous volumes of oil.

Petroleum: Development of this source has been extensive and known resources are being depleted in the U.S. As of 1976, forty-one percent (41%) of all crude oil consumed in the U.S. is now imported from foreign reserves. No crude oil is produced in Cascade County although refining does take place. Crude oil is the mainstay resource of the transportation industry and is being exhausted rapidly. Exploration for new reserves is continuing.

Natural Gas: Cascade County is considered a potential area for natural gas production. Presently, natural gas is supplied by Canadian and Montana fields in the area. Canadian reserves are now being cut off, thus endangering the heating systems for many residential areas. Depletion of this resource can be expected to cause higher prices for consumers. Increased exploration can also be expected.

Wood: Wood is a renewable resource which can be exhausted when demand is too great. There are some excellent forests in the southern part of Cascade County. Inefficient use of this energy resource is its major problem. Ornamental fireplace systems which lose heating potential are very common. Modification of fireplace design would enhance heating systems in many homes.

Renewable sources of energy available to the United States seem to be in the infancy stage of development. Technology to utilize them has often been available for many years but seldom used. Others are still beyond the technology we have today.

Nuclear Fusion: If scientists could successfully harness the fusion reaction, it would provide a nearly unlimited and virtually pollution-free source of power. However, after twenty (20) years of expensive research, they have yet to do this. Some believe the process will never be perfected.

Geothermal Energy: This energy source is one of the few which is environmentally acceptable and yet has the potential of supplying a significant portion of man's energy requirements. Unfortunately, no major hot spots or hot springs exist within Cascade County.

Solar Energy: Today there are no technical barriers to the widespread application of solar energy to meet U.S. needs. Problems exist in that the cost of converting solar heat to useful forms of energy is now higher than conventional sources. Impact upon the environment with this type of development is generally minimal. Small scale projects on an individual basis for home heating are easily developed and can be applicable within Cascade County. Further information on this subject is included later in this appendix.

Wind Energy: Excellent potential exists within Cascade County for this type of development. Community or individual application is possible in many areas of the county. Development of this type of power source is beyond experimental stages for small scale uses. Little affect upon the environment has been observed and the social impacts are also very low.

Fuel from Waste: Production of synthetic gas called methane from waste is possible through solar-organic processes. Implementation of this system would require major changes in waste collection methods. Potential does exist in Cascade County on smaller scales. Adequate sources of sewage, garbage and organic material would be needed thus eliminating some of the pollution problems we now have.

Tide Energy: Potential for this energy source exists in many coastal areas. Several production centers have been constructed but exact impact upon the environment is not known.

Conservation of energy and development of pollution free households can also help alleviate the energy problem. Household conservation tips are available from many different private and government agencies. They include such things as energy saving architecture designs, solar heaters, wind generators, insulation methods, building materials, etc. Pollution prevention devices such as the Clivus Multrum sewage system are also becoming popular in that they also conserve water. Public awareness of the fact that energy and money can be saved through conservation methods is becoming evident. Consciousness that we are still dependent upon energy sources soon to be depleted must be realized and measures must be taken by the individual to begin the shift to alternative energy sources.

*In an effort to clarify this statement, some social costs may include the destruction or loss of any area as open space, the elimination of scenic views or the loss of one's home and property.

Environmental costs, on the other hand, may include such things as air, water or noise pollution and loss of wildlife habitat, farm or grazing lands.

ENERGY RESOURCE COST CHART

ENERGY RESOURCE	FINANCIAL	ENVIRONMENTAL	SOCIAL	AGRICULTURAL	HEALTH	DANGER
Nuclear Fission	very high	high	moderate	low	very high	very high
Nuclear Fusion	very high	low	low	low	low	moderate
Coal Gasification and Liquefaction	very high	very high	high	high (if coal is stripped)	low	low
Oil Shale	very high	very high	high	low	high	low
Geothermal	high	low	low	low	low	? (possible quakes)
Solar	high	low	low	low	none	low
Wind	high	low	low	low	none	low
Fuel from Waste	high	none (enhances)	low	none (benefits)	none	low
Petroleum	high	moderate	moderate	low	low	low
Natural Gas	high	moderate	low	low	low	low
Wood	low	high	low	low	none	low
Tide	high	moderate	moderate	none	low	low

SOLAR ENERGY

Solar energy, man's first energy source, was first used for drying agricultural products and the distillation of liquids many centuries before Christ. Little significant interest was shown in solar energy use until the 20th century when solar-powered electrical plants were constructed in Egypt in 1913. Interest again lagged until the early 1970's when energy shortages forced research into energy sources which were of constant supply (i.e. such as solar power). Today solar energy is considered to have the highest potential for use in the present and future and is now economically competitive with other energy sources.

Solar heating systems usually consist of flat plate or concentrating collectors which transfer heat to a carrying medium, either liquid or air. The heat is then ready for conversion to usable energy forms for use or storage, in rock or eutectic salts.

Collectors on the earth's surface are subject to interruptions by local weather and the intermittent nature of the solar cycle. These may be balanced by large storage capacity or some supplementary energy sources, such as wind, methane extraction or the more conventional natural gas and electricity. Efficiency of the solar system depends upon a number of local variables including sun angle, days of sunlight, and duration of sunlight.

Cascade County has the potential for supplying much of its individual needs through the use of solar energy systems. The average number of clear or partly cloudy days from sunrise to sunset at the Great Falls Airport is 188. The 177 cloudy days will usually give some benefit from solar radiation, even though the effectiveness of a collector may be reduced. Solar energy systems may be more or less effective, depending on specific site conditions. One of the most important conditions is sun angle, which is discussed later.

Some of the advantages attached to solar energy are:

- 1) It is a clean, free, inexhaustible source.
- 2) Solar energy does not add to the thermal burden of the earth.
- 3) Construction of a solar energy system is relatively quick and easy.
- 4) It is safe.
- 5) There is a sufficient quantity.

Disadvantages:

- 1) The collectors work only when the sun is shining, i.e. the energy is intermittent and of low intensity.

SUN ANGLE

The sun angle is the distance of the sun from above the horizon at a certain latitude on any day. For example, if the sun is directly overhead, the sun angle is 90° . The equation for figuring sun angle is:

$$\text{Sun angle} = 90^{\circ} - \phi + \alpha$$

ϕ = Latitude

α = Elevation of the sun (degrees latitude the sun is north or south of the equator)

Example for Great Falls, latitude 47.5°N :

summer solstice (longest day of year) $\alpha = +23.5^{\circ}$ (sun's rays are vertical at 23.5°N)

$$\begin{aligned}\text{Sun angle} &= 90^{\circ} - \phi + \alpha & \phi &= +47.5^{\circ} \text{ (latitude of Great Falls)} \\ &= 90^{\circ} - 47.5^{\circ} + 23.5^{\circ} \\ &= \underline{66^{\circ}} \text{ above horizon}\end{aligned}$$

winter solstice (shortest day of year) $\alpha = -23.5^{\circ}$ (sun's rays are vertical at 23.5°S)


$$\begin{aligned}\text{Sun angle} &= 90^{\circ} - \phi + \alpha & \phi &= +47.5^{\circ} \text{ (latitude of Great Falls)} \\ &= 90^{\circ} - 47.5^{\circ} - 23.5^{\circ} \\ &= \underline{19^{\circ}} \text{ above horizon}\end{aligned}$$

autumnal or vernal equinox (fall or spring, when length of day equals the length of night)

$\alpha = 0^{\circ}$ (sun's rays are vertical at 0° - equator)

$$\begin{aligned}\text{Sun angle} &= 90^{\circ} - \phi + \alpha & \phi &= +47.5^{\circ} \text{ (latitude of Great Falls)} \\ &= 90^{\circ} - 47.5^{\circ} + 0^{\circ} \\ &= \underline{42.5^{\circ}} \text{ above horizon}\end{aligned}$$


Therefore, the sun angle at Great Falls (latitude 47.5°N) varies from 19° on December 22 to 66° on June 22. With this knowledge, it is relatively easy to set up a solar energy system that would make maximum use of the sun's radiation.



WINTER SOLSTICE
DEC. 22



VERNAL & AUTUMNAL EQUINOX
MAR. 23 & SEPT. 22



SUMMER SOLSTICE
JUNE 21

SUN ANGLES at GREAT FALLS, MONT.

LAT. 47.5° N. 12:00 NOON

WIND

Utilizing the power of the atmosphere is not "new"; interest is only reborn. For many, many years wind moved the sailing ships of the world's greatest seapowers. Wind power bore the commerce when this country was young. By 1850 the use of windmills were equivalent to 1.4 billion horsepower hours of work (approximately 1.04 billion Kilowatt hours), but it wasn't until 1894 when the Arctic explorer Narsen used wind to generate electricity.

The Cascade County area has excellent potential for harnessing wind power because of the 13.9 mph average wind velocity found here. Using the December, 1975 National Geographic's example: "Assume that the wind in Chicago blows constantly at (an) average speed of 11.2 miles per hour, but that the average for Minneapolis, also 11.2 mph, comes from winds that blow 8.2 mph for half each day and 14.2 mph for the other half." Because wind power increases as the cube of wind velocity "the city of Minneapolis would have available during its windiest half of the day as much energy as Chicago would have during the whole day." Average wind speed in Great Falls varies with the seasons.

Advantages of using wind:

- 1) Wind is everywhere and free.
- 2) There is no damage to the surrounding environment.
- 3) It does not add to the thermal burden of the Earth.

Disadvantages:

- 1) Windmills may be aesthetically displeasing.
- 2) There are variations in power-plant output because of the fluctuation in duration and intensity of the wind, which necessitates storage facilities.
- 3) To be usable (in most cases) wind speeds must be greater than 7 mph.


ANNUAL ALL WEATHER WIND ROSE

SHOWING PERCENTAGE DISTRIBUTION

GREAT FALLS INTERNATIONAL AIRPORT
GREAT FALLS, MONTANA

PERIOD JAN 1, 1964 - DEC 31, 1967
TOTAL OBS ALL WEATHER: 11,688

PERCENTAGES SHOWN ARE
OF WIND OBS. DURING
ALL WEATHER



Meteorological Data For The Current Year

[illegible]

Normals, Means, And Extremes

Temperatures °F				Normal Dewies days Base 65 °F		Precipitation in inches				Relative humidity pct.				Wind				Mean number of days				Average an on pressure mb	
Normal		Extremes		Normal		Extremes		Normal		Extremes		Normal		Extremes		Normal		Extremes		Normal			
Month	Day	Month	Day	Month	Day	Month	Day	Month	Day	Month	Day	Month	Day	Month	Day	Month	Day	Month	Day	Month	Day		
Jan	20.3	1.9	20.5	1.5	15	1.5	15	1.5	15	1.5	15	1.5	15	1.5	15	1.5	15	1.5	15	1.5	15		
Feb	23.9	1.1	20.5	1.5	15	1.5	15	1.5	15	1.5	15	1.5	15	1.5	15	1.5	15	1.5	15	1.5	15		
Mar	35.9	17.2	26.6	4.0	16.8	2.4	19.69	13.80	0	0.88	2.32	19.65	13.80	0	0.75	2.16	19.67	13.80	0	0.75	2.16		
Apr	40.4	20.6	30.5	7.2	18.6	2.4	19.62	10.79	0	0.87	2.18	19.67	10.79	0	0.87	2.18	19.67	10.79	0	0.87	2.18		
May	54.5	32.3	43.4	8.3	16.9	0.6	19.67	6.48	0	1.18	4.93	19.73	6.48	0	1.18	4.93	19.73	6.48	0	1.18	4.93		
Jun	65.0	43.4	53.3	9.0	17.7	0.6	19.67	3.67	0	2.37	8.13	19.53	3.67	0	2.37	8.13	19.53	3.67	0	2.37	8.13		
Jul	72.1	49.5	60.8	9.7	17.74	3.2	19.69	1.62	3.6	3.11	5.37	19.85	1.62	3.6	3.11	5.37	19.85	1.62	3.6	3.11	5.37		
Aug	83.7	54.9	68.3	10.5	17.6	4.0	19.72	1.8	15.1	1.57	2.32	19.55	1.8	15.1	1.57	2.32	19.55	1.8	15.1	1.57	2.32		
Sep	81.8	53.0	67.1	10.5	17.6	2.3	19.65	2.60	1.6	1.77	4.16	19.74	2.60	1.6	1.77	4.16	19.74	2.60	1.6	1.77	4.16		
Oct	70.0	44.6	57.3	9.8	17.6	3.3	19.72	5.24	2.7	0.68	3.43	19.75	5.24	2.7	0.68	3.43	19.75	5.24	2.7	0.68	3.43		
Nov	59.4	37.1	48.3	8.5	16.64	3.3	19.72	5.24	2.7	0.68	3.43	19.75	5.24	2.7	0.68	3.43	19.75	5.24	2.7	0.68	3.43		
Dec	43.7	34.6	34.6	15.1	19.70	9.12	0	0	0.81	2.27	19.55	0.02	19.54	0	0.81	2.27	19.55	0.02	19.54	0	0.81		
Jan	34.7	18.2	26.5	6.1	16.62	4.3	19.68	11.94	0	0.71	1.73	19.45	11.94	0	0.71	1.73	19.45	11.94	0	0.71	1.73		
Feb	55.9	33.8	44.9	10.6	16.69	4.3	19.68	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13		
Mar	88.7	55.9	68.3	10.6	16.69	4.3	19.68	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13		
Apr	88.7	55.9	68.3	10	16.69	4.3	19.68	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13		
May	88.7	55.9	68.3	10	16.69	4.3	19.68	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13		
Jun	88.7	55.9	68.3	10	16.69	4.3	19.68	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13		
Jul	88.7	55.9	68.3	10	16.69	4.3	19.68	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13		
Aug	88.7	55.9	68.3	10	16.69	4.3	19.68	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13		
Sep	88.7	55.9	68.3	10	16.69	4.3	19.68	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13		
Oct	88.7	55.9	68.3	10	16.69	4.3	19.68	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13		
Nov	88.7	55.9	68.3	10	16.69	4.3	19.68	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13		
Dec	88.7	55.9	68.3	10	16.69	4.3	19.68	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13	19.53	7.852	33.9	14.99	0.13		

Means and extremes above are from existing and comparable exposures. Annual extremes have been exceeded at other sites in the locality as follows: Highest temperature 107 in July 1933; lowest temperature -49 in February 1936; maximum precipitation in 24 hours 3.36 in June 1907.

(a) Length of record, years, through the date of an extreme - The most noted, based on January data.
 (b) 70° and above at Alaskan stations.
 * Less than one half.
 † Trace.

Local Climatological Data

Annual Summary With Comparative Data

1975

GREAT FALLS, MONTANA



Narrative Climatological Summary

The city of Great Falls is located astride the main stem of the Missouri River at its confluence with the Sun River, while the Weather Service Office is located at the Municipal Airport on a plateau between the Sun and Missouri Rivers. This plateau is about 200 feet higher than most of the immediate valley area, and the airport is about two miles southwest of the Sun and Missouri River Junction. Except to the north and northeast, the valley is encircled by mountain ranges, which lie about 30 miles away from east to south, 40 miles to the southwest, and 60 to 100 miles distant from west to northwest. Topography plays an important part in Great Falls' climate. The Continental Divide to the west, and Big and Little Belt Ranges to the south, are primary factors in producing the frequent wintertime "chinook" winds observed in this part of Montana. The valley-plateau combination in the immediate area contributes quite often to marked temperature differences between airport and City proper, either on calm, clear mornings, or when "chinook" winds appear at the airport before they are felt at the lower elevations in town.

Summertime in the area generally is quite pleasant, with cool nights, moderately warm and sunny days, and very little weather that can be called hot or humid. Most summer rainfall occurs in showers or thundershowers, but steady rains do occur during late spring and summer. On the average, the maximum temperature reaches 100° or better during four summers out of 10 and an average year will have 20 days with maximums 90° or higher. Freezing temperatures do not occur in July or August, very rarely in June, and are observed only on two or three days in the usual May or September. Frost occurs frequently in April and October, but more often in the valleys than on the surrounding hills or plateau.

Winters are not so cold as is usually expected of a continental location at this latitude, largely as a result of the "chinook" winds for which this area is noted. While sub-zero weather is experienced normally several times during a winter, the coldest weather seldom lasts more than a few days at a time, and is usually terminated by southwest "chinook" winds which can produce sharp temperature rises of 40° or more in 24 hours. As a result of recurring "chinooks" throughout the winter season, snow seldom lies on the ground for more than a few days. In fact, the ground usually is bare, or nearly bare of snow, most of the winter, except in the surrounding mountains and higher foothills. On the other hand, invasions of cold air from the polar regions occur a few times each winter, and sharp temperature falls from above freezing to below zero within 24 hours are observed occasionally from mid-December to March.

Precipitation generally falls as snow during late fall, winter, and early spring, although rain can occur in any month. Late spring, summer and early fall precipitation is almost always rain, but some hail is observed occasionally during summer thundershowers.

Although Great Falls' average annual precipitation would normally classify the area as semi-arid, it is important to note that about 70 percent of the annual total falls normally during the April-September growing season. The combination of ideal temperatures during the peak of the growing season, long hours of summer sunshine, and nearly 10 inches of precipitation during the six critical months, makes the climate very favorable for dryland farming. Heavy fog seldom occurs, incidence usually being limited to about one day per month, but each case lasts only a small part of the day. Although the average windspeed is relatively high, extremely strong winds (over 70 mph) seldom are observed, and visibility normally is excellent (15 miles or more).

Average Temperature

Year	Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec	Annual
1936	19.8	-5.2	34.0	45.0	63.0	66.4	76.4	70.2	58.8	51.3	34.0	25.4	45.3
1937	3.1	17.8	34.0	46.5	57.4	62.3	72.6	88.0	58.4	52.7	34.0	26.0	44.4
1938	31.0	18.2	34.2	45.8	52.1	61.9	69.3	67.2	65.5	50.2	33.0	29.3	46.5
1939	32.6	18.7	34.0	48.4	57.7	57.0	72.8	69.6	59.9	47.6	45.2	36.6	48.5
1940	15.8	22.7	34.6	41.1	57.4	64.6	71.3	70.4	62.4	51.6	27.2	32.8	46.2
1941	28.6	28.4	34.4	44.4	55.5	62.4	71.3	67.6	51.8	46.2	39.5	28.5	46.6
1942	27.6	20.0	31.1	47.0	50.4	56.8	69.3	67.2	57.2	50.1	33.3	25.2	45.0
1943	11.7	30.1	24.6	49.6	50.4	56.6	69.4	68.1	59.2	52.1	39.3	32.8	45.4
1944	33.3	24.0	24.9	47.2	57.2	57.6	67.6	64.8	59.4	55.4	37.6	28.6	46.2
1945	28.6	23.6	31.7	38.8	51.1	56.8	71.7	70.2	54.6	51.6	33.4	24.0	45.0
1946	31.6	32.1	41.6	50.0	51.0	61.4	70.6	68.7	57.0	40.5	24.1	26.8	46.4
1947	25.8	23.2	27.4	46.0	55.1	58.3	73.2	68.6	56.6	51.7	30.2	31.0	45.4
1948	29.3	19.9	27.2	43.2	53.0	61.3	68.4	68.6	60.0	50.6	35.6	18.4	44.5
1949	10.0	14.6	31.2	51.0	56.8	61.2	68.6	71.0	58.4	41.5	47.5	19.8	44.7
1950	-3.6	35.3	24.9	40.7	53.8	58.5	66.3	64.5	55.8	49.2	31.7	33.0	42.5
1951	14.2	23.7	19.1	40.8	53.1	54.1	68.2	62.7	57.6	47.0	34.5	15.2	40.5
1952	18.3	27.5	24.0	50.9	54.8	61.2	66.5	66.3	61.3	50.2	33.6	22.5	45.9
1953	22.4	31.2	34.0	39.2	49.1	58.5	70.0	69.1	60.9	54.2	44.8	32.5	48.1
1954	10.2	36.6	24.9	37.0	53.5	56.8	71.5	65.1	55.8	49.8	44.7	34.6	45.0
1955	25.9	21.6	24.9	40.5	49.5	60.4	67.0	70.3	56.0	50.4	19.4	22.3	42.2
1956	20.8	23.3	33.8	41.4	54.3	63.8	69.4	65.1	58.8	47.7	36.3	29.2	45.5
1957	7.9	22.4	31.8	42.8	56.4	61.9	71.8	65.6	59.3	40.7	30.0	36.6	44.5
1958	36.6	23.6	24.1	44.0	61.9	58.9	63.8	71.2	58.5	52.2	33.3	29.9	46.8
1959	24.1	16.9	38.1	43.4	48.5	63.0	70.8	65.8	55.7	44.8	29.7	36.0	44.7
1960	22.8	23.9	31.3	42.8	52.3	62.2	74.5	65.5	60.2	50.2	35.5	30.2	46.0
1961	32.8	36.0	37.7	40.3	54.8	69.4	70.2	72.4	50.2	46.6	30.0	20.3	46.7
1962	19.0	20.7	24.9	48.1	50.6	61.4	65.0	67.3	58.7	50.5	40.3	32.6	46.8
1963	12.8	36.8	39.4	42.9	52.9	61.4	66.9	68.6	64.0	54.1	38.6	23.5	47.0
1964	28.7	33.0	26.9	42.3	51.8	62.5	72.4	65.4	53.9	53.0	31.9	11.2	44.7
1965	23.9	27.0	20.9	44.7	51.4	60.3	69.6	68.0	45.0	53.4	24.9	29.1	46.0
1966	13.9	27.1	34.4	40.2	55.8	59.9	70.0	65.1	64.2	47.4	29.0	27.9	44.8
1967	26.3	32.1	27.2	35.5	52.5	60.2	73.0	71.0	62.9	50.2	34.2	22.1	45.8
1968	22.0	33.2	40.4	40.6	49.5	59.2	67.8	64.4	56.5	48.0	35.8	16.1	44.5
1969	-2.8	16.6	26.2	50.3	55.8	59.3	68.3	72.1	61.3	38.1	40.1	29.9	42.9
1970	14.7	37.3	29.0	38.5	53.8	66.5	71.4	70.9	54.6	44.7	30.2	23.0	44.1
1971	16.2	29.4	31.6	40.5	54.6	62.0	67.5	76.0	54.7	44.7	36.3	18.0	44.6
1972	12.8	22.5	38.3	42.4	53.5	63.2	64.7	69.4	53.9	43.2	26.3	17.9	43.4
1973	24.9	29.6	39.4	40.2	55.4	63.6	71.4	71.2	58.2	49.2	25.2	28.9	46.4
1974	19.8	33.8	33.3	47.1	49.5	66.9	72.9	62.9	54.8	51.4	38.7	32.5	47.0
1975	22.7	13.1	27.4	30.9	50.0	58.8	71.8	62.9	57.3	45.7	32.9	28.6	42.0
RECORD	21.2	26.1	31.4	43.3	53.3	60.9	69.7	67.9	57.6	45.3	34.8	27.1	45.1
MEAN	30.4	35.9	41.4	54.3	64.9	72.4	83.8	81.8	69.9	59.2	43.8	35.8	56.1
MAX	12.0	16.3	21.3	32.2	41.7	49.3	55.5	53.9	45.2	37.4	25.6	18.4	34.1

Precipitation

Year	Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec	Annual
1936	0.59	1.20 ¹	0.43	0.71	1.18	2.12	0.19	0.43	1.17	0.34	0.15	0.56	9.18
1937	0.55	0.16	0.51	0.22	0.27	3.29	1.43	0.61	1.17	0.33	0.43	0.78	9.85
1938	0.03	0.56	0.44	0.27	3.72	4.52	1.00	1.64	0.75	1.55	0.32	0.02	16.84
1939	0.29	0.33	0.24	0.88	1.52	2.41	0.66	0.59	1.30	1.21	0.15	0.31	9.89
1940	0.62	0.69	0.83	1.78	2.10	0.91	0.73	0.05	2.24	0.83	1.13	1.16	12.09
1941	0.29	0.36	0.55	1.34	2.07	3.67	2.27	0.88	3.58	0.55	1.11	1.19	17.64
1942	0.46	1.02	0.84	0.55	4.64	2.44	1.50	0.67	1.06	0.56	0.94	0.06	14.55
1943	1.48	0.62	0.51	1.26	1.03	5.32	1.01	0.84	0.61	0.57	0.39	0.42	13.96
1944	1.44	1.47	0.75	1.15	3.68	1.24	1.44	1.31	0.04	0.99	0.50	1.21	16.21
1945	0.32	0.53	0.76	0.67	1.49	3.24	0.20	0.85	2.60	0.80	0.32	1.73	13.51
1946	0.10	0.12	0.42	0.48	1.70	1.96	1.92	1.18	1.93	1.53	1.78	0.86	13.98
1947	0.50	0.71	1.47	0.40	0.68	3.42	0.26	0.30	2.52	0.54	1.11	0.44	12.35
1948	1.23	0.36	1.42	0.61	3.43	4.84	2.28	0.50	0.72	0.08	0.39	0.53	16.39
1949	1.40	0.79	1.11	0.41	3.21	1.54	0.94	0.27	0.52	0.90	0.05	0.53	11.67
1950	0.91	0.01	2.14	1.47	0.67	2.76	3.54	2.94	0.31	0.03	1.19	0.58	16.55
1951	0.41	1.34	1.27	2.50	2.23	3.57	2.54	2.50	1.44	1.72	0.41	1.71	21.59
1952	0.24	1.63	0.76	0.30	1.57	1.22	0.15	1.21	0.24	0.15	1.36	0.09	9.02
1953	0.55	1.46	0.74	1.85	8.13	5.06	0.06	0.56	1.14	0.09	0.13	0.98	20.75
1954	1.23	0.21	1.01	0.86	1.17	4.71	0.63	2.63	1.86	1.35	0.02	1.7	15.68
1955	0.80	1.47	1.53	2.48	4.26	2.23	4.32	0.04	0.16	0.54	2.27	0.55	19.65
1956	0.52	0.37	0.38	0.55	1.33	2.58	1.02	1.68	0.44	1.02	0.30	0.59	10.76
1957	1.60	0.73	0.77	0.95	2.82	2.94	0.75	1.11	1.68	1.89	0.61	0.11	16.18
1958	0.55	2.16	1.24	0.67	1.09	4.68	2.32	0.42	0.28	0.31	1.24	1.17	16.14
1959	1.57	1.00	0.58	0.36	2.94	1.88	0.04	0.27	1.55	1.20	1.71	0.43	13.63
1960	0.28	0.52	0.15	2.13	1.71	0.52	0.39	2.66	0.43	0.04	0.79	0.19	9.61
1961	0.22	0.19	0.84	0.94	1.80	0.73	1.01	0.63	1.95	0.32	1.49	0.30	10.48
1962	1.29	0.95	0.74	0.58	1.18	2.30	1.09	1.89	1.10	1.17	0.36	0.51	19.95
1963	1.71	0.32	0.35	1.24	1.27	2.68	0.96	0.49	0.87	0.63	0.21	1.02	11.95
1964	0.65	0.32	1.74	1.91	3.26	4.34	1.50	1.66	0.28	1.7	0.72	1.23	17.91
1965	0.84	1.18	0.79	2.51	1.47	5.27	1.03	1.58	1.90	1.7	1.13	0.59	18.39
1966	1.63	0.51	0.79	0.74	1.54	2.17	1.81	0.77	1.62	1.32	1.62	0.99	14.10
1967	1.12	0.25	2.14	3.69	2.17	3.65	0.91	0.23	1.59	1.13	0.26	1.47	18.68
1968	1.29	0.24	0.90	1.11	2.64	2.69	0.06	2.16	2.92	0.11	0.68	1.38	16.36
1969	2.05	0.40	0.44	0.38	1.14	5.33	1.11	0.03	0.13	0.59	0.11	0.40	12.41
1970	0.99	1.02	1.14	1.85	3.16	2.32	1.16	0.77	0.67	1.00	0.53	0.70	15.34
1971	1.22	0.65	1.12	0.66	3.03	0.62	0.27	1.16	0.61	0.30	0.36	1.48	11.48
1972	1.47	0.62	1.01	0.77	1.59	0.94	1.51	1.26	0.85	1.17	0.20	1.68	13.07
1973	0.33	0.26	0.30	2.69	0.95	1.43	0.13	0.68	1.29	0.97	1.38	1.37	12.16
1974	1.44	0.26	1.10	1.03	3.16	1.08	0.48	4.76	0.73	0.36	0.26	0.60	15.26
1975	1.14	0.71	1.34	4.63	3.89	4.47	1.20	2.13	0.74	3.43	1.01	0.55	25.24
RECORD	0.87	0.70	0.91	1.27	2.40	2.89	1.16	1.19	1.14	0.80	0.76	0.72	14.82

* Indicates station move or relocation of instruments. See Station Location table.

Record values are shown through the current year for the period beginning in 1936. Data are from Cooperative locations through March 1937 and from Airport locations thereafter.

Heating Degree Days

GREAT FALLS, MT

Season	July	Aug	Sept	Oct	Nov	Dec	Jan	Feb	Mar	Apr	May	June	Total
1945-56	88	8	300	444	1364	1318	1363	1198	960	700	341	94	4178
1946-57	40	77	203	537	790	1107	1770	1189	962	681	259	124	7714
1947-58	7	69	197	750	694	877	874	1153	1138	623	144	194	5920
1948-59	100	13	235	400	944	1081	1297	1347	828	637	505	109	7502
1949-50	26	67	310	621	1055	890	1301	1187	1041	660	409	114	7481
1950-61	11	76	187	454	878	1073	989	807	839	732	326	14	6389</

STATION LOCATION

GREAT FALLS, MONTANA

Location	Occupied from	Occupied to	Airline distance and direction from previous location	Latitude North	Longitude West	Elevation above										Remarks
						Sea level	Ground								Sea level	
							Ground at ice-pressure site	Wind instruments	Extreme thermometers	Psychrometer	Telepsychrometer	Tipping bucket rain gage	Weighing rain gage	8" rain gage		
COOPERATIVE																
End of Central Avenue Across Park Drive	11 01 91	6 30 98		47° 0'	111° 18'	3331		4					5			
Corner Central Avenue & Second Street	7 01 98	2	500 ft E	47° 30'	111° 18'	3330		25					18			9 - Sometime after 3 22 06, but before 7 15 13.
Post Office Grounds 1st Avenue N & 2nd Street	*	10 16 14	600 ft NE	47° 31'	111° 18'	3328		4					3			
423 Fourth Avenue North	00 17 14	01 18	1400 ft NE	47° 1'	111° 18'	3334		4					3			
1709 Third Avenue North	4 01 18	9 30 19	5500 ft E	47° 31'	111° 17'	3387		4					3			
412 Thirteenth Street North, Fire Station	00 01 14	31 37	2100 ft NE	47° 1'	111° 18'	3370		4					3			
AIRPORT																
Municipal Airport Gore Field, Mail Hangar 3 1/2 miles SW of P. O.	11 01 31	12 19 39		47° 29'	111° 22'	3654	35	4	1				3			SAKRS station to 11/1/36, then CAA.
Municipal Airport Gore Field, Administration Building	12 20 39	8 01 59	No Change	47° 29'	111° 22'	3664	b75	a18	a17		c15	c15	15	c 3697		Weather Bureau from 1 25 40. a - 16 feet 2 22/40 to 11 21 41 b - 63 feet to 6/30/42. c - Added 11/21/41.
International Airport Administration Building	8 01 59	Present	No Change	47° 29'	111° 22'	e3662	f22	18	17		15	15	15	d4 3692	d	d - Commissioned 2 4 61 1100 feet W of thermometer site. e - 3664 feet to 2 4 61. f - 23 feet to 11 10 64. g - Decommissioned May 1975.

Requests for additional climat. information should be addressed to: Director, National Climatic Center, Federal Building, Asheville, N. C. 28801

Sale Price: 20 cents per copy. Checks and money orders should be made payable to Department of Commerce, NOAA. Remittances and correspondence regarding this publication should be sent to: National Climatic Center, Federal Building, Asheville, N. C. 28801. Attn: Publications.

I certify that this is an official publication of the National Oceanic and Atmospheric Administration, and is compiled from records on file at the National Climatic Center, Asheville, North Carolina 28801.

Thomas D. Potter
 Director, National Climatic Center

USCOMM-NOAA-ASHEVILLE - 900

U.S. DEPARTMENT OF COMMERCE
 NATIONAL CLIMATIC CENTER
 FEDERAL BUILDING
 ASHEVILLE, N.C. 28801

AN EQUAL OPPORTUNITY EMPLOYER

POSTAGE AND FEES PAID
 U.S. DEPARTMENT OF COMMERCE

210



FIRST CLASS



For more information:

- GPO - Catalog on Solar Energy Heating and Cooling Products
(ERDA-75_ stock #052-010-00470-1 \$3.80)
- TIC - Solar Energy Program (ERDA-15)
pamphlet length, general description of ERDA's solar program
- GPO - National program for Solar Heating and Cooling
(Residential and Commerical Applications) (ERDA-23A)
stock #052-020-00475-1 \$2.50
- GPO - Definition Report, National Solar Energy Research, Development and
Demonstration Program (ERDA-49)
ERDA research plans for photovoltaic, wind, biomass, ocean and solar
thermal conversion applications, as well as for heating and cooling
stock #052-020-00473-5 \$2.00
- NTIS - Summary Report, Federal Wind Energy Program (ERDA-84)
overview of research and development activities in wind energy,
with abstracts of individual projects. \$5.00
- GPO - Solar Energy Utilization for Heating and Cooling (NSF 74-41)
Currently available design data
stock #3800-00188 \$.70
- NTIS - Solar Energy: A Bibliography (TID-3351-R1)
Bibliographic citations of scientific and technical inforamtion on
solar energy in the ERDA energy information data base through 1975
Part I (citations) - \$13.75 Part II (index) - \$10.75
- TIC - Solar Energy Update
A monthly abstract-index journal covering research, development, and
demonstration information from ERDA and its contractors, with reference
to reports from other agencies, institutes, and countries.
 - Annual Cycle Energy Systems (ACES)
Pamphlet-length description of one energy conservation demonstration
project. Available from: Division of Building and Industries, ERDA,
Washington, D.C. 20545
- TIC - Solar energy reprints from ERDA
Pocket folder of articles for the layman reprinted from Popular Science,
Sunset, etc.
- TIC - Solar Energy
Popular-level pamphlet (EDM-527) and more detailed booklet (IB-801)
describing solar energy applications

TIC - I've got a question about using solar energy. (EDM-816)
 Answers to the questions people ask ERDA about the solar demonstration power and the homeowner's use of solar energy.

OEO - Save Energy: Save Money!
 OEO pamphlet 6143-5
 Simple, low-cost methods of insulation, using heat, and utilizing the sun.

FEA - Tips for Energy Savers

ECR - The Energy Crisis -- What can we do?
 \$1.00

SHC - Living with the Energy Crisis
 8 - page brochure \$.25

CP - 350 Ways to Save Energy and Money in Your Home and Car
 by Spies, Konzo, Calvin, and Thenis
 198 page paperback book \$3.95

UNH - How to Set Up a Coal or Wood Stove
 a 2 - page brochure

UNH - Home Heating in an Emergency
 15 page booklet

UP - Save Your Heating Dollars -- and Keep Your Home Warm
 a series of 1 - page brochures

ASE - Spectrum - An Alternative Technology Equipment Directory
 \$1.00

CM - A waterless, non-chemical method of individual sewage treatment

J - Wood and coal heaters

SAB - Energy Earth and Everyone: A Global Strategy for Spaceship Earth
 by Medard Gabel \$4.95

For information marked:

OEO - Office of Economic Opportunity

Regional Energy Coordinator

Denver Regional Office - VIII

Federal Building

1961 Stout Street

Denver, Colorado 82002

TIC - ERDA Technical Information Center

P.O. Box 62

Oak Ridge, Tennessee 37830

NTIS- National Technical Information Service

5285 Port Royal Road

Springfield, Virginia 22161

GPO - Superintendent of Documents

US Government Printing Office

Washington, DC 20402

ASE - Alternative Sources of Energy

Route 2, Box 90A

Milaca, Minnesota 56353

CM - Glenn Nelson

Route 2, Meadows Road OR

Whitefish, Montana 59937

J - Jøtul

Tim Dowell OR

1703 McMannamy Draw

Kalispell, Montana 59901

Clivus Multrum USA

14 - A Eliot Street

Cambridge, Massachusetts 02138

Importers:

Kristia Associates

Box 1118

Portland, Maine 04104

SAB - Straight Arrow Books

625 Third Street

San Francisco, California 94107

FEA - Office of Public Affairs

Federal Energy Administration

Washington, DC 20461

ECR - Energy Conservation Research

9 - Birch Road

Malvern, Pennsylvania 19355

SHC - Small Homes Council

University of Illinois

Champaign, Illinois 61820

CP - Crown Publishing Company

New York, New York

UNH - Cooperative Extension Service

University of New Hampshire

Durham, New Hampshire 03824

UP - Cooperative Extension Service

University of Pennsylvania

University Park, Pennsylvania 16802

GLOSSARY

GLOSSARY

ADEL MOUNTAIN VOLCANICS - 3,200 feet of potash-rich volcanic and related intrusive rocks (Tertiary).

AESTHETIC - Pertaining to a feeling or sense of the beautiful.

ALLUVIAL FAN - A fan- or cone-shaped deposit of alluvium formed where a stream flows out onto a level area or into a slower stream.

ALLUVIUM - Material, including sand, clay, gravel, and mud, deposited in river beds, lakes, alluvial fans, valleys and elsewhere by modern streams.

BASALT - A dark colored igneous rock.

BEDROCK - The solid rock beneath the soil and subsoil.

BENCHES - A name applied to ledges of all kinds of rock that are shaped like steps or terraces.

BENTONITE - A clay, formed from the decomposition of volcanic ash, that absorbs water readily.

BITUMINOUS COAL - A mineral that burns with a yellow smoky flame; soft coal.

BLACKLEAF FORMATION - Very black to pink, green and lavender shale and light colored fine-grained quartz sandstone. Much bentonite. (Lower Cretaceous)

BLOWOUT - An area from which soil material has been removed by wind. Such an area appears as a nearly barren, shallow depression with a flat or irregular floor consisting of a resistant layer, an accumulation of pebbles, or wet soil lying just above a water table.

BOOTLEGGER MEMBER (of Blackleaf Formation) - Interbedded gray sandstone, siltstone and shale, and dark gray shale. (Lower Cretaceous)

CARRYING CAPACITY - The capability of a particular environment to support animal life.

CITIZEN'S ADVISORY COMMITTEE (CAC) - A body made up of residents from within a planning district who will make recommendations on planning related matters to the Planning Board.

CLAY - 1. Mineral soil material that is plastic when wet, consisting essentially of hydrated silicates of aluminum. 2. Soil material that contains 40 percent or more of clay, less than 45 percent of sand, and less than 40 percent of silt.

CLAY LOAM - Soil material that contains 27 to 40 percent of clay and 20 to 45 percent of sand.

COLLUVIUM - Mixed deposits of soil material and rock fragments accumulated near the base of steep slopes through soil creep, slides, and local wash.

COLORADO SHALES - Dark gray to gray-black shale. Some siltstone and sandstone and limestone beds locally. (Upper and Lower Cretaceous)

CONGLOMERATE - Rock composed of rounded pebbles cemented together in a matrix of finer material.

CRETACEOUS PERIOD - From 135 million years before present to 70 million years before present.

DEVELOPMENT - Any change in land-use that raises the economic value of a parcel of land.

DIASTROPHISM - All the processes that change the shape of the earth's surface, producing mountains and valleys, continents and ocean basins, etc.

DIKE - 1. A relatively thin, flat surface mass of igneous rock that cuts across the structure of the enclosing rocks or extends through massive rocks. Usually formed by the intrusion of magma into a fissure. 2. An embankment, usually of natural materials.

DISTURBED BELT - The sedimentary layers disturbed during the uplift of the Rocky Mountains.

FAULT - A fracture in soil or in a rock mass along which movement has occurred, causing one side to be displaced in relation to the other.

FERDIG-KEVIN MEMBERS - Gray to dark gray shale, some concretions, numerous bentonite beds. Members of Marias River Shale Formations. (Upper Cretaceous)

FLOOD - A temporary rise in the level of water which results in an inundation of areas not ordinarily covered by water.

100 year flood - A flood of a magnitude such that it has a 1% chance of occurring in any given year.

5 year flood - A flood of a magnitude such that it has a 20% chance of occurring in any given year.

10 year flood - A flood of a magnitude such that it has a 10% chance of occurring in any given year.

20 year flood - A flood of a magnitude such that it has a 5% chance of occurring in any given year.

FLOOD MEMBER - Brown, fine-grained, flaggy to massive, ledge-making sandstone with thin interbeds of dark gray shale. Member of the Blackleaf Formation. (Lower Cretaceous)

FLOODPLAIN - A plain built up by stream deposition which may be submerged periodically by floodwaters.

FLOODWAY - The normal stream channel and that area adjoining the floodplain needed to convey moving waters at flood stage.

FLOODWAY FRINGE - The area of the floodplain lying outside the floodway, but subject to periodic inundation from flooding.

FLOWEREE-CONE MEMBERS - Dark gray and very dark gray shales. (Upper Cretaceous)

FOLD - A bend in rock layers caused by compression of the earth's crust.

FORMATION - A group of strata or layers having certain common characteristics that more or less readily distinguish them from adjacent layers.

GROUNDWATER - All water below the land surface. It has its origin in downward seepage of surface water to a layer of impervious material.

IGNEOUS ROCK - Rock formed by the solidification of magma.

INTEGRITY - An unimpaired condition.

INTRUSION - A body of igneous rock that has penetrated older rock.

INTRUSIVE ROCKS - Rocks that have solidified at some distance beneath the surface of the earth.

JURASSIC PERIOD - From 180, million years before present to 135, million years before present.

KOOTENAI FORMATION - Three (3) or more sandstone bodies with interbedded vari-colored claystone and siltstone. (Lower Cretaceous)

LANDSLIDE - The downward movement of large masses of rocks and earth.

LIMESTONE - A sedimentary rock composed mostly of calcium carbonate.

LOAM - The textural class name for soil having a moderate amount of sand, silt, and clay. Loam soils contain 7 to 27 percent of clay, 28 to 50 percent of silt, and less than 52 percent of sand.

LOAMY SANDS - (see sandy loam.)

MAGMA - Molten or otherwise fluid rock occurring naturally within the earth. Igneous rocks are believed to have been formed by solidification of magma.

MARIAS RIVER FORMATION - Dark gray shale divided into four (4) members.

METAMORPHIC ROCK - A rock that has been greatly altered from its previous condition through the combined action of heat and pressure. For example, marble is a metamorphic rock produced from limestone, gneiss is produced from granite, and slate is produced from shale.

MISSION CANYON FORMATION - Mostly massive brown to gray limestone. (Upper Kinderhook to Osage)

MORRISON FORMATION - Green, drab, or gray marl containing in lower two-thirds (2/3) numerous lenticular bodies of limestone. Near the base is a persistent sandstone unit. Upper one-third (1/3) contains a preponderance of sandstone beds. (Upper Jurassic)

PERFORMANCE STANDARDS - Regulations involving a specific quantitative measurement of an effect of impact of a particular activity.

PYRITE - Fool's gold. It is an iron and a source of sulfur, but is not used as an iron source because of the difficulty of extracting the mineral from the ore.

QUATERNARY PERIOD - From about 2 million years before present.

REVERSE FAULT - An earth fault in which one block has been pushed up and over the other.

ROCK FALL - The process by which rock becomes detached from a nearly vertical slope resulting in a talus slope.

SAND - Individual rock or mineral fragments in soils having diameters from 0.5 to 2.0 millimeters. Usually sand grains consist mainly of quartz but they may be of any mineral composition. The textural class name of any soil that contains 85 percent or more of sand and not more than 10 percent of clay.

SANDSTONE - A porous sedimentary rock consisting of sand held together by such materials as silica or limestone.

SANDY LOAM - Soil of textural class which generally contains 50 percent sand and less than 20 percent clay.

SCARP - An escarpment, which is the steep-faced edge of a cuesta or plateau.

SEDIMENTARY ROCK - A rock composed of particles deposited from suspension in water. Chief groups of sedimentary rocks are conglomerates, from gravels; sandstones, from sand; shales, from clay; and limestones, from soft masses of calcium carbonate. There are many intermediate types. Some wind-deposited sands have been consolidated into sandstones.

SHALE - A laminated, fine-grained sedimentary rock derived from clay muds.

SILL - An intrusion of igneous rock approximately uniform in thickness, much wider than it is thick, and approximately parallel to the bedding of the sedimentary rock in which it has been emplaced.

SILT - 1. Individual mineral particles of soil that range in diameter between the upper size of clay, 0.002 millimeter and lower size of very fine sand, 0.05 millimeter. Soil of the textural class of silt contains 80 percent or more of silt and less than 12 percent of clay. 2. Sediments deposited from water in which the individual grains are approximately the size of silt, although the term is sometimes applied loosely to sediments containing considerable sand and clay.

SILT LOAM - Soil material having (a) 50 percent or more of silt and 12 to 27 percent of clay or (b) 50 to 80 percent of silt and less than 12 percent of clay.

SILT-SHALES - Shales composed of silt-sized particles.

SILTSTONE - A very fine-grained sandstone, the particles of which are predominantly of silt grade.

SILTY CLAY - Soil of this textural class has 40 percent or more clay and 40 percent or more of silt.

SILTY CLAY LOAM - Soil of this textural class has 27 to 40 percent clay and less than 20 percent sand.

SLUMPING - The process by which a more or less continuous mass of earth slips downhill along a curved basal slip plane in such a way that the mass of earth rotates backwards, toward the slope. This is usually a relatively rapid movement.

SOIL CREEP - Slow movement of rock fragments downslope, occurring most commonly when lower soil is nearly saturated with water.

STRATUM (pl. Strata) - A layer of sedimentary rock, or a group of layers consisting throughout of approximately the same material. Stratum, bed, and layer are approximately equivalent terms.

SUBSOIL - A common term that can not be defined accurately. It has been carried over from early days when "soil" was conceived only as the plowed soil and that under it as the "subsoil."

SWIFT FORMATION - Dark gray non-calcareous shale overlain by fine-grained glauconitic sandstone. (Upper Jurassic)

TAFT HILL-VAUGHN MEMBERS - Dull gray, greenish to pinkish bentonitic siltstone. Members of Blackleaf Formation. (Lower Cretaceous)

TALUS - A mass of rock debris that collects at the bottom of a cliff.

TECTONIC PLATE - Any one of the huge movable segments into which the earth's crust is divided.

TELEGRAPH CREEK FORMATION - Yellow (outcrop) to gray-white (subsurface), partly shaly sandstone and sandy shale and gray shale. Usually thin bedded. (Upper Cretaceous)

TERRACE - A relatively flat earth surface, usually long and narrow, bounded by an ascending slope on one side and a descending slope on the other.

TERTIARY PERIOD - From 70, million years before present to about 2, million years before present.

VIRGELLE SANDSTONE MEMBER - Buff to yellow (outcrop) gray to white (subsurface) fine- to coarse-grained, often salt-and-pepper sandstone. Often forms massive cliffs. Member of Eagle Sandstone. (Upper Cretaceous)

VOLCANICS - Those igneous rocks which have reached or nearly reached the surface before solidifying, the type of which is a lava flow.

WIND EROSION - The removal, transportation, and deposition of rock material by wind.

BIBLIOGRAPHY

PARTIAL BIBLIOGRAPHY

- Alden, William C., Physiography and Glacial Geology of Eastern Montana and Adjacent Areas, USGS Professional Paper 174, Government Printing Office, Washington, D.C., 1932.
- Allen, Thomas B., Vanishing Wildlife of North America, National Geographic Society, Washington, D.C., 1974.
- Alt, David D. and Donald W. Hyndman, Roadside Geology of the Northern Rockies, Mountain Press Publishing Company, Missoula, Montana, 1972.
- Balster, Clifford A., editor, Catalog of Stratigraphic Names for Montana, State of Montana Bureau of Mines and Geology, Butte, Montana, 1971.
- Cascade County: County Profiles, Montana Department of Intergovernmental Relations, Division of Research and Information Systems, Helena, Montana, 1974.
- Cascade County Situation Statement 1972, Prepared by Cascade County USDA Committee for Rural Development.
- Durrenberger, Robert W., Dictionary of the Environmental Sciences, National Press Books, Palo Alto, California, 1973.
- Environmental Plan for Northeast Clatsop County, Oregon, Prepared by the Clatsop County Planning Department, Astoria, Oregon, 1975.
- Environmental Plan for Southeast Clatsop County, Oregon, Prepared by the Clatsop County Planning Department, Astoria, Oregon, 1975.
- Environmental Plan for the Clatsop Plains, Prepared jointly by the Clatsop County Planning Department, Citizen Advisory Committees, and OSU Extension Service, September, 1974.
- Fisher, Cassius A., Geology and Water Resources of the Great Falls Region Montana, USGS Water Supply Paper 221, Government Printing Office, Washington, D.C., 1909.
- Gabel, Medard, Energy Earth and Everyone: A Global Strategy for Spaceship Earth, Straight Arrow Books, San Francisco, California, 1975.
- General Soil Map and Interpretations for Resource Planning in Cascade County, Montana, SCS, 1969.

Hamilton, Roger "Can We Harness the Wind?" National Geographic, Washington, D.C., December, 1975, pp. 812 - 829.

Ice Jams on the Missouri River, communication with William Murray, Coordinator, Cascade County Civil Defense, July, 1976.

I've got a question about using solar energy, Energy Research and Development Administration, Washington, D.C., May, 1976.

Local Climatological Data for Great Falls, Montana, 1975, NOAA Environmental Data Service, Asheville, North Carolina.

McKey, Jeff and Mike Morgan, An Environmental Plan of the Lewis & Clark, Youngs, and Walluski River Valleys, Western Interstate Commission for Higher Education, Boulder, Colorado, 1974.

Morgan, Mike and Jeff McKey, An Environmental Plan of Southeast Clatsop County, Western Interstate Commission for Higher Education, Boulder, Colorado, 1974.

1970 Census of Population and Housing Census Tracts: Great Falls, Montana, Standard Metropolitan Statistical Area, U.S. Bureau of Census, Government Printing Office, Washington, D.C., 1971.

Performance Standards . . . a technique for controlling land use, OSU Extension Service, Special Report 424, Corvallis, Oregon, 1974.

Rare and Endangered Species, Communication with Montana Fish and Game Commission, 1976.

Rice, C. M. Dictionary of Geological terms, Edwards Brothers, Inc., Ann Arbor, Michigan, 1945.

Small, Cooley and Associates, Cascade County, Montana: Comprehensive Plan for Sewer and Water, Great Falls, Montana, 1969.

Soil Survey Reconnaissance: Central Montana, SCS with Montana Agricultural Experiment Station, Series 1940 Number 9, Washington, D.C., February, 1953.

Solar Energy and Conservation for Home Heating and Cooling, Energy Research and Development Administration, Washington, D.C., May, 1976.

Thurow, Charles, William Toner and Duncan Erley, Performance Controls for Sensitive Lands: A Practical Guide for Local Administrators, ASPO Planning Advisory Service, Chicago, Illinois, 1975.

Water Resources Survey: Cascade County, Montana, State Engineer's Office, Helena, Montana, June, 1961.

This intern report was read and accepted by a staff member at:

Agency: Cascade County Planning Board

Address: County Courthouse
Great Falls MT 59401

This report was completed by a WICHE intern. This intern's project was part of the Resources Development Internship Program administered by the Western Interstate Commission for Higher Education (WICHE).

The purpose of the internship program is to bring organizations involved in community and economic development, environmental problems and the humanities together with institutions of higher education and their students in the West for the benefit of all.

For these organizations, the intern program provides the problem-solving talents of student manpower while making the resources of universities and colleges more available. For institutions of higher education, the program provides relevant field education for their students while building their capacity for problem-solving.

WICHE is an organization in the West uniquely suited for sponsoring such a program. It is an interstate agency formed by the thirteen western states for the specific purpose of relating the resources of higher education to the needs of western citizens. WICHE has been concerned with a broad range of community needs in the West for some time, insofar as they bear directly on the well-being of western peoples and the future of higher education in the West. WICHE feels that the internship program is one method for meeting its obligations within the thirteen western states. In its efforts to achieve these objectives, WICHE appreciates having received the generous support and assistance of the Economic Development Administration; the Jessie Smith Noyes Foundation; the National Endowment for the Humanities; the Wyoming Office of Manpower Planning; and of innumerable local leaders and community organizations, including the agency that sponsored this intern project.

For further information, write Bob Hullinghorst, Director, Resources Development Internship Program, WICHE, P.O. Drawer 'P', Boulder, Colorado 80302 or call (303) 492-7177.

8421451000045800:
1.1C:1076:WICHE:2H330

